

**G-protein mediated inhibitory effect of a nitric oxide donor
on the L-type Ca²⁺ current in rat ventricular myocytes**

Najah Abi-Gerges #, Rodolphe Fischmeister, Pierre-François Méry

*Laboratoire de Cardiologie Cellulaire & Moléculaire, INSERM U-446,
Faculté de Pharmacie, Université Paris-Sud, F-92296 Châtenay-Malabry, France*

Short title: Inhibition of cardiac $I_{Ca,L}$ by NO donors

Key words: L-type Ca²⁺ channel, NO donors, G protein, isoprenaline, guanylyl cyclase, cyclic AMP, cyclic GMP

#Corresponding author:

Dr. Najah ABI-GERGES
INSERM U-446
Faculté de Pharmacie
F-92296 Châtenay-Malabry Cedex
France

Tel.: (33-1) 46 83 57 57

Fax: (33-1) 46 83 54 75

E-mail: Naja.Abi-g@cep.u-psud.fr

Abstract

- 1- The role of the cGMP pathway in the modulation of the cardiac L-type Ca^{2+} current ($I_{\text{Ca,L}}$) by nitric-oxide (NO) was examined in rat ventricular myocytes.
- 2- The NO-donors DEANO, SIN-1, SNP, SNAP and GSNO had no significant effects on basal $I_{\text{Ca,L}}$. However, DEANO (100 μM) inhibited $I_{\text{Ca,L}}$ after the current had been previously stimulated by either isoprenaline (Iso, 1-10 nM), a β -adrenergic agonist, or isobutylmethyl-xanthine (IBMX, 10-80 μM), a wide spectrum phosphodiesterase (PDE) inhibitor.
- 3- The anti-adrenergic effect of DEANO on $I_{\text{Ca,L}}$ was not mimicked by other NO-donors (SIN-1, SNAP and SPNO).
- 4- The NO-sensitive guanylyl cyclase inhibitor, ODQ (10 μM), antagonized the inhibitory effect of DEANO on $I_{\text{Ca,L}}$. Likewise, inhibitors of the cGMP-dependent protein kinase (cG-PK), Rp-8-chloro-phenylthio-cGMP (10 μM) and KT5823 (0.1 and 0.3 μM), also abolished the inhibitory effect of DEANO on Iso (1-10 nM)-stimulated $I_{\text{Ca,L}}$.
- 5- Intracellular dialysis with exogenous cAMP (10-100 μM) blunted the inhibitory effect of DEANO (10 and 100 μM) on $I_{\text{Ca,L}}$. SNAP and SNP also had no effect on the cAMP-stimulated $I_{\text{Ca,L}}$.
- 6- Pre-treatment of the myocytes with Pertussis toxin (0.5 $\mu\text{g/ml}$, 4-6 hours at 37°C) eliminated the inhibitory effect of DEANO (100 μM) on $I_{\text{Ca,L}}$, in the presence of either Iso (0.01 and 1 nM) or IBMX (10-80 μM).
- 7- These results demonstrate that DEA-NO produces anti-adrenergic effects in rat ventricular myocytes. This effect of DEA-NO occurs in a cGMP-dependent manner, and involves activation of cG-PK and regulation of a Pertussis toxin-sensitive G-protein.

INTRODUCTION

The synthesis of nitric oxide (NO) plays an important role in the endothelium-dependent relaxation of various blood vessels. In the heart, the role of NO synthesis is rather controversial (reviewed in Méry *et al.* 1997; Feron *et al.* 1999; Kojda & Kottenberg, 1999). NO was soon recognised as an activator of cGMP synthesis in the whole heart, as well as in purified cardiac myocytes. In turn, cGMP co-ordinates the activity of various proteins, such as the cGMP-stimulated phosphodiesterase (PDE2), the cGMP-inhibited-PDE (PDE3), and the cGMP-dependent protein kinase (cG-PK). Accordingly, several NO-donors can modulate myocyte shortening, ionic currents, and contractile proteins, in a cGMP-dependent manner. More recently, nitrosothiols (SNAP, SNAC, GSNO) or the sydnonimine SIN-1 were found to elicit cGMP-independent effects in cardiac myocytes, including the modulation of calcium channels (Hu *et al.* 1997; Xu *et al.* 1998), the inhibition of the creatine kinase (Gross *et al.* 1996), and the modulation of mitochondrial respiration (Wolin *et al.* 1997).

While these results tend to support a physiological role of NO in heart muscle, a number of questions remain unanswered. First, in several studies performed on isolated cardiac myocytes, NO-donors had no effect on cell shortening (Stein *et al.* 1993; McDonnell *et al.* 1995; 1997), L-type Ca^{2+} current ($I_{\text{Ca,L}}$) (Thomas *et al.* 1997), or the cAMP-activated chloride current (Zakharov *et al.* 1996). Thus, subtle differences between cardiac preparations can blunt the effects of NO at the single cell level. Second, the exact mechanism by which NO or cGMP produces its effect in a given cardiac preparation is not clear. For instance, the inhibitory effect of NO or cGMP on $I_{\text{Ca,L}}$ in mammalian cardiac myocytes may take place either via activation of PDE2 (Feron *et al.* 1999) or activation of cG-PK (Méry *et al.* 1991; Sumii & Sperelakis, 1995; Whaler & Dollinger, 1995). Third, while the effect of cG-PK on $I_{\text{Ca,L}}$ appears to take place at the level of the L-type Ca^{2+} channel (or a closely associated

protein Méry *et al.* 1991; Sumii & Sperelakis, 1995), the possibility exists that cG-PK might also act upstream from Ca²⁺ channel phosphorylation. Indeed, in smooth muscle cells, cG-PK was shown to directly phosphorylate α_i subunits of GTP-binding proteins (G proteins) and/or receptors (Pfeiffer *et al.* 1995; Wang *et al.* 1998). In addition, cG-PK can increase the spontaneous binding of GTP on α subunits and reduce the stimulatory effects of receptor agonists on their GTPase activity (Pfeiffer *et al.* 1995; Wang *et al.* 1998; Miyamoto *et al.* 1997).

In the present study, we examined the effects of different NO-donors on basal and stimulated $I_{Ca,L}$ in ventricular myocytes isolated from rat hearts. In this preparation, we found earlier that cG-PK mediates the inhibitory effect of exogenous cGMP on $I_{Ca,L}$ (Méry *et al.* 1991). While none of the NO-donors tested produced any effect on basal $I_{Ca,L}$, the NONOate DEANO produced a profound inhibition of the current stimulated by isoprenaline. We show that cG-PK mediates this effect through the activation of a Pertussis toxin-sensitive G protein.

Part of this work has been presented in an abstract form (Abi-Gerges *et al.* 1997c).

METHODS

The investigation conforms with our institution guidelines which are defined by the European Community guiding principles in the care and use of animals (86/609/CEE, *CE Off J* n°L358, 18 December 1986) and the French decree n°87/848 of October 19, 1987 (*J Off République Française*, 20 October 1987, pp. 12245-12248). Authorizations to perform animal experiments according to this decree were obtained from the French Ministère de l'Agriculture et de la Forêt (n°04226, April 12, 1991).

Cell isolation and storage

Male Wistar rats (160-250 g) were anaesthetised by intraperitoneal injection of urethane (2g/kg) and heparin (2.5mg/kg). After all reflex activity had ceased, the animal was killed by opening the chest and rapidly removing the heart. Myocytes were dispersed using collagenase A (0.255mg/ml; Boehringer-Mannheim Biochemica, Mannheim, Germany) as previously described (Scamp *et al.* 1990; Abi-Gerges *et al.* 1997a).

Electrophysiology

The whole-cell configuration of the patch-clamp technique was used to record the L-type Ca^{2+} current ($I_{\text{Ca,L}}$) on Ca^{2+} -tolerant cells, as described (Scamps *et al.* 1990; Abi-Gerges *et al.* 1997a; 1999). The routine protocol consisted in a depolarizing pulse to 0 mV test potential (400 ms duration) elicited every 8 sec from a holding potential of -50 mV. The test potential was set at 0 mV because, at this potential, the $I_{\text{Ca,L}}$ amplitude (Scamps *et al.* 1990; Méry *et al.* 1991; Abi-Gerges *et al.* 1999 ; see Fig. 2) and steady-state activation are at their maximal values in rat myocytes (Scamps *et al.* 1990). Current-voltage relationships and inactivation curves were performed as described (Abi-Gerges *et al.* 1999). The experiments were carried at room temperature (22-32°C, mean value $25.6 \pm 0.1^\circ\text{C}$, n=346), and the temperature did not change by $>2^\circ\text{C}$ in any given experiment.

During patch-clamp experiments, the maximal amplitude of $I_{Ca,L}$ and the current at the end of the 400 ms test potential were measured as described (Abi-Gerges *et al.* 1997a). Currents were not compensated for capacitive and leak currents. The density of basal $I_{Ca,L}$ was 5.25 ± 0.14 pA pF⁻¹ and the density of the current at the end of the 400 ms pulse was 0.37 ± 0.02 pA pF⁻¹ (n=266). The steady-state value of the end-pulse current was stable over the time-course of the experiments (see figures). The effects of the agonists used in this study were not correlated with the amplitude of the end-pulse current (data not shown). The decay of the capacitive transient was fast (< 3 ms), and did not interfere significantly with the activation of the calcium current (mean time to peak = 6.2 ± 0.1 ms, n=266). On-line analysis of the recordings was made possible by programming a PC-compatible 486/50 microcomputer in Assembly language (Borland) to determine, for each membrane depolarization, peak and steady state current values.

Solutions for patch-clamp recordings

The extracellular solution contained in (mM): 107 NaCl, 10 HEPES, 20 CsCl, 4 NaHCO₃, 0.8 NaH₂PO₄, 1.8 MgCl₂, 1.8 CaCl₂, 5 D-glucose, 5 Na-pyruvate, and 6×10^{-4} tetrodotoxin, pH 7.4 adjusted with CsOH (294 mosm). Solutions were superfused onto floating myocytes as described (Abi-Gerges *et al.* 1997a). The patch pipettes (0.5-1.0 MΩ) were filled with an intracellular solution containing in (mM): 119.8 CsCl, 5 EGTA (acid form), 4 MgCl₂, 5 Na₂-phosphocreatine, 3.1 Na₂ATP, 0.42 Na₂GTP, 0.062 CaCl₂ (pCa 8.5) and 10 HEPES, pH 7.3 adjusted with CsOH (292 mosm).

Drugs

3-Morpholiniosydnonimine (SIN-1) was a generous gift from Doctor J. Winicki (Hoechst-Houdé Laboratories, Paris-La Défense). Spermine-NONOate (SPNO) and S-Nitroso-L-glutathione (GSNO) were from Alexis Corp. (La Jolla, CA, USA); S-Nitroso-N-acetyl-D,L-

penicillamine (SNAP), 2-(N,N-Diethylamino)-diazolate-2-oxide (DEANO), 1H-[1, 2, 4]oxadiazolo[4, 3-a]quinoxaline-1-one (ODQ) from Tocris-Cookson (Bristol, UK) or Alexis Corp.; 8-(4-Chlorophenylthio)guanosine-3'-5'-cyclic monophosphorothioate, Rp-isomer (Rp8cG) from Biolog L.S.I. (Bremen, Germany); 8-(4-Chlorophenylthio)-guanosine-3',5'-cyclic monophosphate (8-p-CPT-cGMP) and KT5823 from Calbiochem-France Biochem (Meudon, France); Bordetella Pertussis toxin from Sigma-Aldrich (Saint Quentin Fallavier, France); Tetrodotoxin from Latoxan (Rosans, France). All others drugs were from Sigma-Aldrich. Drugs were prepared and used according to manufacturer's instructions. Mocked-DEANO consisted in a DEANO (100 μ M)-containing solution left at room temperature for >20h. Solutions were prepared by dilution to the desired concentration in the physiological solution at the beginning of each experiment.

Statistical analysis

Results are expressed as mean \pm S.E.M. Differences between mean values were tested for statistical significance by paired Student's *t*-test, as indicated. In the text, the "basal" condition for $I_{Ca,L}$ refers to the absence of extracellular isoprenaline or IBMX, or intracellular cAMP.

RESULTS

Effect of NO-donors on basal $I_{Ca,L}$

NO-donors can modulate basal $I_{Ca,L}$ in some, but not all cardiac myocytes (Kirstein *et al.* 1995; Wang *et al.* 1998; Campbell *et al.* 1996; Méry *et al.* 1993). In the experiment of Fig. 1A, a rat myocyte was first exposed to control intracellular and extracellular solutions. Under this condition, the amplitude of the basal $I_{Ca,L}$ declined slowly ($\sim -5.4 \text{ pA min}^{-1}$), a phenomenon known as “run-down”. Superfusion of the myocyte with DEANO (100 μM) barely modified the amplitude and the kinetics of the basal $I_{Ca,L}$. As summarized in Fig. 1B, the basal $I_{Ca,L}$ amplitude was not significantly modified by different NO-donors. These included the NONOate DEANO (100 μM), the nitrosothiols GSNO (1 mM) and SNAP (1 mM), the ferrocyanate SNP (500 μM), and the sydnonimine SIN-1 (1 mM). Lower concentrations of these compounds (down to 1 μM) also had no effect on basal $I_{Ca,L}$ (data not shown).

(Figure 1 near here)

Since nitrosylation and/or oxidation might account for effects of NO-donors on the basal $I_{Ca,L}$ in other preparations (Campbell *et al.* 1996; Hu *et al.* 1997) we further investigated the effects of reducing or oxidant agents sensitivity on the basal $I_{Ca,L}$ in rat myocytes. The basal $I_{Ca,L}$ was not changed when the myocyte was superfused with 1 mM reduced glutathione ($5.9 \pm 2.6 \%$ over basal, $n=5$), 1 mM DL-DTT ($2.3 \pm 1.6 \%$ over basal, $n=6$), 0.1 mM N-acetylpenicillamine ($-1.6 \pm 0.9 \%$ over basal, $n=4$). In contrast, extracellular application of the superoxide anion generator LY 83583 (10 μM) inhibited the basal $I_{Ca,L}$ ($-36.9 \pm 2.5 \%$ over basal, $n=17$, $p<0.001$). Therefore, the basal activity of L-type Ca^{2+} channels in rat myocytes was sensitive to an oxidative treatment but not to NO-donors.

Inhibitory effect of DEANO on the β -adrenergic stimulation of $I_{Ca,L}$

NO-donors can reduce the β -adrenergic stimulation of $I_{Ca,L}$ in frog and guinea-pig ventricular myocytes (Méry *et al.* 1993; Levi *et al.* 1994; Whaler & Dollinger, 1995; reviewed in Méry *et al.* 1997). We first investigated the effects of DEANO in the presence of isoprenaline (Iso), a non-selective β -adrenergic agonist. In the experiment of Fig. 2A, a myocyte was exposed to control solutions and the $I_{Ca,L}$ amplitude reached a steady-state 11.8 min after the beginning of the recording of $I_{Ca,L}$. The myocyte was then superfused with 10 nM Iso, which induced a ~2-fold increase in $I_{Ca,L}$. In the continuing presence of Iso, superfusion with 100 μ M DEANO elicited a pronounced reduction in the response of $I_{Ca,L}$ to Iso. The inhibitory effect of DEANO was reversible. Figure 2B shows the current-voltage relationships of $I_{Ca,L}$ (*filled symbols*) and of the steady-state current measured at the end (400 ms) of the test pulse (*empty symbols*) in the same experiment as in Fig. 2A. The effects of Iso and DEANO on $I_{Ca,L}$ were homogenous along the voltage range and neither drug produced a change in the steady-state current (similar findings were obtained in 10 other experiments). In addition, DEANO did not change the inactivation curve of $I_{Ca,L}$ (data not shown). Thus, DEANO produced an anti-adrenergic effect on $I_{Ca,L}$ which occurred in a voltage-independent manner.

(Figure 2 near here)

The inhibitory effect of DEANO (100 μ M) was studied at three different Iso concentrations (1, 3 and 10 nM). As summarized in Fig. 3, increasing the Iso concentration progressively reduced the inhibitory effect of DEANO on $I_{Ca,L}$ suggesting that the inhibitory effect of DEANO was somewhat competitive with the stimulatory effect of Iso. The involvement of NO in the anti-adrenergic effect of DEANO was investigated using mocked-DEANO (see

Methods). The Iso (1 nM)-stimulation of $I_{Ca,L}$ was identical in the absence and in the presence of 100 μ M mocked DEANO (respectively, $176.8 \pm 11.9\%$ and $188.9 \pm 10.1\%$ of control, $n=4$). In addition, the stimulatory effect of 1 nM Iso on $I_{Ca,L}$ was not changed in the presence of 100 μ M diethylamine (respectively $179.1 \pm 16.0\%$ and $177.0 \pm 20.0\%$ of control, $n=4$), the other metabolite of DEANO. Thus, the inhibitory effect of DEANO on Iso-stimulated $I_{Ca,L}$ was mediated by NO and not by one of its metabolites.

(Figure 3 near here)

Effect of other NO-donors on the β -adrenergic stimulation of $I_{Ca,L}$

For a comparison with the effect of DEANO, we examined the effects of other NO-donors on the Iso-stimulated $I_{Ca,L}$, namely SPNO (a NONOate like DEANO), SIN-1 (a sydnonimine) and SNAP (a nitrosothiol). Fig. 4A shows a typical experiment in which SIN-1 (100 μ M) had no effect on $I_{Ca,L}$ stimulated by 10 nM Iso. As summarised in Fig. 4B, SIN-1 (100 μ M) as well as the two other NO-donors, SPNO (100 μ M) and SNAP (100 μ M), produced inconsistent and non significant effects on $I_{Ca,L}$ stimulated with either 1 or 10 nM Iso. In addition, the shape of the current-voltage relationship of the Iso-stimulated $I_{Ca,L}$ was not modified in the presence of either SIN-1, SNAP, or SPNO (respectively, $n=8$, 5, and 2, data not shown). These negative results were not due to the washout of some cellular component in the whole-cell configuration of the patch clamp technique since SNAP also failed to modify the Iso-stimulated $I_{Ca,L}$ when tested in the perforated-patch configuration (data not shown, see also Thomas *et al.* 1997). There were not due either to a difference in temperature, since the mean temperatures of the experiments was similar with DEA-NO (25.3 ± 0.3 °C, $n=25$), SPNO (28.0 ± 0.5 °C, $n=5$), SIN-1 (28.5 ± 0.5 °C, $n=3$) and SNAP (28.3 ± 0.4 °C, $n=4$) when tested on top of 1 nM Iso. Thus, among a total of four NO-donors tested, DEANO was the only one

to produce consistently an inhibitory effect on Iso-stimulated $I_{Ca,L}$ in rat ventricular myocytes (see Maragos *et al.* 1991; Feelisch *et al.* 1991).

(Figure 4 near here)

Contribution of guanylyl cyclase to the anti-adrenergic effect of DEANO on $I_{Ca,L}$

To evaluate the contribution of cGMP to the modulation of $I_{Ca,L}$ by DEANO, we examined the effect of DEANO in the presence of ODQ, an inhibitor of the NO-sensitive guanylyl cyclase (Kojda *et al.* 1996; Abi-Gerges *et al.* 1997b). In the typical experiment of Fig. 5A, a myocyte was exposed to control solutions, and then superfused with 3 nM Iso. The stimulatory effect of the β -adrenergic agonist was reduced by about 50% upon addition of DEANO (100 μ M). In the continuing presence of Iso plus DEANO, superfusion with ODQ (10 μ M) increased $I_{Ca,L}$, eliminating most of the inhibitory effect of DEANO. The effect of ODQ was slowly reversible upon washout of the drug. On average (Fig. 5B), ODQ (10 μ M) fully antagonized the inhibitory effects of DEANO on $I_{Ca,L}$, in the presence of either 1, 3 or 10 nM Iso. However, ODQ (10 μ M) did not modify Iso (10 nM)-stimulated $I_{Ca,L}$ in the absence of DEANO ($-2.2 \pm 2.1\%$ over Iso level, $n=4$). Thus, these results indicate that the inhibitory effect of DEANO on Iso-stimulated $I_{Ca,L}$ in rat ventricular myocytes is mediated by activation of the NO-sensitive guanylyl cyclase.

(Figure 5 near here)

Contribution of cG-PK to the anti-adrenergic effect of DEANO on $I_{Ca,L}$

We have shown earlier that an intracellular application of cGMP in rat ventricular myocytes produces an inhibition of cAMP-stimulated $I_{Ca,L}$ that involves the activation of cG-PK (Méry

et al. 1991). Similarly, the Iso (1 nM)-stimulation of $I_{Ca,L}$ was strongly reduced in the presence of 8-p-CPT-cGMP (100 μ M), a selective activator of cG-PK (from $235.1 \pm 8.9\%$ to $168.2 \pm 6.9\%$ stimulation, $n=4$, $p<0.005$). Next, the role of the cG-PK in the anti-adrenergic effect of DEANO was examined using (Rp)-8-CPT-cGMP (Rp8cG), a membrane-permeant inactive analog of cGMP which competes with cGMP for the binding on cG-PK (Butt *et al.* 1994). In the typical experiment of Fig. 6A, the β -adrenergic stimulation of $I_{Ca,L}$ (with 1 nM Iso) was strongly inhibited by DEANO (100 μ M). This anti-adrenergic effect of DEANO was fully antagonized by superfusion of the cell with 10 μ M Rp8cG, in a reversible manner. Fig. 6B summarizes the results of several similar experiments. At 1 and 10 nM Iso, Rp8cG (10 μ M) fully reversed the inhibitory effect of DEANO (100 μ M) on $I_{Ca,L}$. We also examined the effects of KT5823, another cG-PK inhibitor structurally unrelated to cGMP (Komalavila & Lincoln, 1996). As summarized in Fig. 6B, KT5823 (0.1 or 0.3 μ M) also fully reversed the anti-adrenergic effect of DEANO (100 μ M). In the absence of DEANO, neither Rp8cG (10 μ M) nor KT5823 (0.1-0.3 μ M) had any significant effect on Iso-stimulated $I_{Ca,L}$ (data not shown). Thus, these results demonstrate that cG-PK mediates the inhibitory effect of DEANO on $I_{Ca,L}$ in rat ventricular myocytes.

(Figure 6 near here)

Effect of DEANO on cAMP-stimulated $I_{Ca,L}$ in rat ventricular myocytes

When elicited by intracellular dialysis of cGMP in rat ventricular myocytes, cG-PK-dependent inhibition of $I_{Ca,L}$ appeared to take place at the level of the Ca^{2+} channels (Méry *et al.* 1991; Sumii & Sperelakis, 1995). In other cell types, cG-PK can modulate initial steps of signal transduction, at the level of receptors and/or G proteins (Pfeiffer *et al.* 1995; Wang *et al.* 1998, Miyamoto *et al.* 1997). To discriminate between these possibilities, we first studied the

effect of DEANO in myocytes dialysed with cAMP through the patch pipette, in order to bypass the steps involved in cAMP production. In the experiment of Fig. 7A, the patch pipette was filled with a control intracellular solution supplemented with 30 μM cAMP, and intracellular dialysis of the myocyte started when the membrane patch was ruptured. The diffusion of cAMP into the cytosol induced a large increase in $I_{\text{Ca,L}}$ (to 44.9 pA pF^{-1}). Surprisingly, superfusion of the myocyte with DEANO (100 μM) did not modify the cAMP-stimulated $I_{\text{Ca,L}}$. The effect of another NO-donor, SNP, is shown in Fig. 7B. In this experiment, the myocyte was first exposed to control intracellular and extracellular solutions. Intracellular dialysis with cAMP (10 μM) started at the time indicated by the arrow, inducing a 2.5-fold increase in $I_{\text{Ca,L}}$ (to 26.7 pA pF^{-1}). Superfusion of the myocyte with increasing concentrations of SNP (5, 50 and 500 μM) had no effect on the cAMP-stimulated $I_{\text{Ca,L}}$. The effects of DEANO, SNP and SNAP on $I_{\text{Ca,L}}$ in cAMP (10-100 μM) dialyzed myocytes are summarized in Fig. 7C. On average, the density of $I_{\text{Ca,L}}$ was elevated to $23.02 \pm 3.57 \text{ pA pF}^{-1}$ ($n=12$). None of the NO-donors did significantly change the cAMP-stimulated $I_{\text{Ca,L}}$. Thus, since DEANO inhibits the Iso-stimulated $I_{\text{Ca,L}}$ but not the cAMP-stimulated $I_{\text{Ca,L}}$, it must act at a step located upstream from cAMP production in the cAMP signalling cascade leading to $I_{\text{Ca,L}}$ stimulation. In the following, we tested the hypothesis that DEANO via activation of cG-PK interferes with the receptor-dependent modulation of adenylyl cyclase.

(Figure 7 near here)

Effect of DEANO on Iso-stimulated $I_{\text{Ca,L}}$ in Pertussis toxin treated myocytes

In vitro experiments demonstrated that cG-PK could modulate the activity of G_i proteins, which are negatively coupled to adenylyl cyclase in rat cardiac myocytes (Pfeiffer *et al.* 1995; Miyamoto *et al.* 1997). Therefore, we studied the effect of DEANO in Pertussis toxin (PTX)-

treated myocytes, where G_i/G_o proteins are irreversibly inactivated. Myocytes were incubated for 4 to 6 hours (before patch-clamp experiments) in the presence of $0.5 \mu\text{g ml}^{-1}$ PTX, at 37°C (Hilal-Dandan *et al.* 1992). In the typical experiment shown in Fig. 8A, Iso (1 nM) was applied on a PTX-treated myocyte, inducing a fast and large rise in $I_{Ca,L}$. However, under these conditions, superfusion of the myocyte with DEANO (100 μM) had no effect on $I_{Ca,L}$. Since the muscarinic receptor agonist acetylcholine (ACh) produces an anti-adrenergic effect on $I_{Ca,L}$ which clearly involves activation of PTX-sensitive G proteins (reviewed in Méry *et al.* 1997; Feron *et al.* 1999), we tested the efficacy of PTX to inactivate G_i/G_o proteins by investigating the effect of a subsequent application of ACh (1 μM) on Iso-stimulated $I_{Ca,L}$. As shown in Fig. 8A, and summarized in Fig. 8B, PTX-treatment fully abrogated the anti-adrenergic effect of either ACh (1 μM) as well as that of DEANO (100 μM). Note that when normalized to the Iso-stimulated $I_{Ca,L}$ amplitude, ACh, but not DEANO, tended to potentiate the effect of the β -adrenergic agonist in PTX-treated myocytes (Fig. 8C).

We next examined whether the effect of PTX was taking place upstream or downstream from cG-PK activation. For this, we investigated the effect 8-p-CPT-cGMP (100 μM) in PTX-treated myocytes. In four such cells, the stimulation of $I_{Ca,L}$ by Iso 1 nM was only slightly attenuated by 8-p-CPT-cGMP (from $192.0 \pm 11.0\%$ to $171.7 \pm 11.0\%$ of control, $p=0.1$), demonstrating that PTX-treatment blunted the inhibitory effect of a direct activation of the cG-PK on $I_{Ca,L}$. In these cells, the Iso-stimulated $I_{Ca,L}$ was also not affected by ACh (1 μM , $198.9 \pm 8.0\%$ of control). Altogether, these data demonstrate that DEANO inhibits $I_{Ca,L}$ via a cG-PK modulation of a PTX-sensitive G_i/G_o protein.

(Figure 8 near here)

Effect of DEANO on IBMX-stimulated $I_{Ca,L}$ in Pertussis toxin treated myocytes

While PTX treatment of rat ventricular myocytes did not modify basal $I_{Ca,L}$ density (5.2 ± 0.4 pA pF⁻¹, n=23, in PTX vs. 5.3 ± 0.1 pA pF⁻¹, n=266, in control), it somewhat increased the Iso response (at 0.01 and 1 nM, compare Fig. 8B and Fig. 3). Thus, the anti-adrenergic effect of DEANO might have been blunted in PTX-treated-myocytes as a result of a saturating production of cAMP. Therefore, we re-examined the involvement of G_i/G_o proteins in the modulation of $I_{Ca,L}$ by DEANO under conditions where intracellular cAMP was non maximally increased. To do this, we used low concentrations of isobutyl-methyl-xanthine (IBMX), a broad spectrum phosphodiesterase inhibitor which, unlike Iso, does not activate cAMP synthesis but increases cAMP level by reducing its degradation. Figure 9A shows a typical experiment performed in a control (untreated) myocyte. IBMX (40 μM) elicited a 25% increase in basal $I_{Ca,L}$ and this effect was totally abolished by further application of DEANO (100 μM). This effect of DEANO clearly involved NO generation, since the addition of the NO-scavenger carboxy PTIO (Akaike *et al.* 1993) totally antagonized the effect of the NO-donor. Indeed, in three myocytes where the stimulation of $I_{Ca,L}$ by 40 μM IBMX (154.2 ± 4.9 % of control) had been reduced by 100 μM DEANO (to 118.5 ± 2.9 % of control, p<0.005), the addition of carboxy PTIO (100 μM) to the DEANO solution restored the stimulation of $I_{Ca,L}$ to its level in Iso alone (to 156.8 ± 1.7 % of control).

In contrast to control cells, DEANO failed to inhibit the response of $I_{Ca,L}$ to IBMX in PTX-treated myocytes. In the typical experiment of Fig. 9B, neither DEANO (100 μM) nor ACh (1 μM) produced any inhibition of $I_{Ca,L}$ stimulated by IBMX (20 μM). The results of several similar experiments are summarized in Fig. 9C. On average, IBMX (10-80 μM) exerted submaximal stimulatory effects on $I_{Ca,L}$. While ACh (1 μM) strongly reduced the IBMX response in control (untreated) myocytes, this inhibition was absent in PTX-treated myocytes indicating that G_i/G_o proteins were efficiently blocked by PTX. Similarly, DEANO (100 μM)

strongly antagonized the IBMX-stimulated $I_{Ca,L}$ in untreated myocytes, and the effect was abolished in PTX-treated myocytes. Thus, the inhibitory effect of DEANO on $I_{Ca,L}$ shared some similarity with the effect of ACh, in that both effects required the integrity of PTX-sensitive G proteins. Note that in one out of the nine experiments in PTX-treated myocytes, ACh had no effect on IBMX-stimulated $I_{Ca,L}$ while DEANO still elicited an inhibitory effect.

(Figure 9 near here)

DISCUSSION

In this study, we report that $I_{Ca,L}$ is regulated by NO in rat ventricular cardiac myocytes. The NO-donor DEANO exerted a pronounced anti-adrenergic effect, which involved the activation of the NO-sensitive guanylyl cyclase and cG-PK. The main locus of action of cG-PK appeared to be a PTX-sensitive G protein.

In agreement with studies in frog and guinea-pig ventricular myocytes (Levi *et al.* 1994; Whaler & Dollinger, 1995; Méry *et al.* 1993), we found that a NO-donor, DEANO, strongly inhibited the β -adrenergic stimulation of $I_{Ca,L}$ in rat ventricular myocytes. The inhibitory effect of DEANO occurred in the micromolar range of concentrations and was not mimicked by mocked-DEANO or diethylamine. This effect clearly involved NO generation, since the addition of the NO-scavenger carboxy PTIO (Akaike *et al.* 1993) totally antagonized the effect of the NO-donor. In contrast, other NO-donors (SIN-1, SNAP and SPNO, a NONOate like DEANO) did not significantly modify the Iso-stimulated $I_{Ca,L}$. However, the rate of NO release by these NO-donors is slow compared to that of DEANO (Feelisch *et al.* 1991; Maragos *et al.* 1991; Ferrero *et al.* 1999). As suggested by Kojda *et al.* (1996), the time course of NO release appears critical when comparing the functional effects of different NO-donors. Nevertheless, although DEANO is 1000-fold more potent than SIN-1, SNP or SNAP, all these NO-donors inhibit the Iso-stimulated $I_{Ca,L}$ in frog ventricular myocytes (*e.g.* see Méry *et al.* 1993; Abi-Gerges *et al.* 1997b). A possible explanation for this discrepancy may come from the observation that rat and mouse ventricles exhibit higher myoglobin contents than other species (O'Brien *et al.* 1992). Since myoglobin is known as a potent NO scavenger, it may hinder the effects of slow NO sources, such as SIN-1, SNAP and SPNO, but not that of a fast NO source like DEANO (Beckman & Koppenol, 1996; Ishibashi *et al.* 1992).

Another interesting observation was that DEANO behaved as an apparent competitive antagonist of the response of $I_{Ca,L}$ to Iso in rat myocytes. We found that the inhibitory effect of DEANO on $I_{Ca,L}$ was reduced as the concentration of Iso was increased from 1 to 10 nM. This observation might help to explain the discrepancy between negative and positive results obtained in the literature when comparing the efficacy of a given NO-donor in different cardiac preparations (Thomas *et al.* 1997).

DEANO was shown to enhance cGMP production in isolated rat ventricular myocytes (Kojda *et al.* 1996). In agreement with this finding, we found that activation of the cGMP pathway accounted for the anti-adrenergic effect of DEANO on $I_{Ca,L}$ in the same preparation. Indeed, the effect of DEANO on $I_{Ca,L}$ was antagonized by ODQ, a specific guanylyl cyclase inhibitor (Abi-Gerges *et al.* 1997b; Sandirasegarane & Diamond, 1999). Moreover, the effect of DEANO on $I_{Ca,L}$ was mimicked by 8-p-CPT-cGMP, a cG-PK activator, and antagonized by Rp8cG and KT5823, two cG-PK inhibitors (Butt *et al.* 1994; Komalavila & Lincoln, 1996). Therefore, the endogenous activation of the cG-PK accounted for the anti-adrenergic effect of DEANO on $I_{Ca,L}$.

Earlier studies in rat ventricular myocytes have shown that cG-PK was involved in the inhibitory effects of exogenous cGMP on $I_{Ca,L}$ (Méry *et al.* 1991; Sumii & Sperelakis, 1995). Intracellular dialysis of isolated myocytes with cGMP or with constitutively active cG-PK elicited inhibitory effects on $I_{Ca,L}$ when the current was maximally stimulated with 100 μ M IBMX or 100 μ M intracellular cAMP. In addition, NO-donors inhibited IBMX- or cAMP-stimulated $I_{Ca,L}$ in guinea-pig myocytes (Levi *et al.* 1994; Whaler & Dollinger, 1995). Accordingly, L-type Ca^{2+} channels were viewed as a major target of cG-PK in these cells (Méry *et al.* 1991; Levi *et al.* 1994; Whaler & Dollinger, 1995; Sumii & Sperelakis, 1995).

However, we found here that in rat ventricular myocytes the inhibitory effect of DEANO on $I_{Ca,L}$ was strongly reduced when the concentration of Iso was increased to 10 nM (Fig. 3) or in the presence of 200 μ M IBMX (data not shown). Moreover, DEANO did not inhibit the stimulation of $I_{Ca,L}$ induced by intracellular cAMP. Therefore, the major target of cG-PK may differ when activated by endogenous cGMP production (this study) or by intracellular dialysis of exogenous cGMP (Méry *et al.* 1991). Our present results suggest that cG-PK acts at the level of G proteins when guanylyl cyclase is stimulated by NO since the inhibitory effect of DEANO on $I_{Ca,L}$ was antagonized by PTX, which inactivates G_i/G_o proteins. Both G_i and G_o proteins are involved in the regulation of $I_{Ca,L}$ (Méry *et al.* 1997; Feron *et al.* 1999; Valenzuela *et al.* 1997), but only G_i -dependent pathways were shown to be modulated by cG-PK (Pfeiffer *et al.* 1995). Activation of cG-PK was shown to phosphorylate α_i subunits of G proteins (Pfeiffer *et al.* 1995) and to increase the spontaneous binding of GTP (Miyamoto *et al.* 1997). Our results strongly suggest that this modulation of α_i subunits of G_i proteins accounts for the observed cG-PK-dependent, PTX-sensitive inhibitory effect of DEANO on Iso-stimulated $I_{Ca,L}$ in rat myocytes. Yet, we can only speculate on the contribution of this mechanism in the regulation of the Iso-stimulated $I_{Ca,L}$ by NO in guinea-pig myocytes (Levi *et al.* 1994; Whaler & Dollinger, 1995). Interestingly, rat ventricles express higher levels of myoglobin than guinea-pig ventricles (O'Brien *et al.* 1992), and high myoglobin content blunted cGMP production induced by NO (Ishibashi *et al.* 1992). Therefore, we propose that only high levels of cGMP can elicit the direct inhibition of the L-type calcium channels by the cG-PK (as seen during dialysis of cGMP, or in guinea-pig myocytes), while moderate levels of cGMP induce the specific regulation of G_i proteins by cG-PK (as seen in rat myocytes).

In addition to the regulation of cGMP production, NO can regulate directly the activity of several proteins in the heart, including the L-type Ca^{2+} channels. Nitrosothiols and SIN-1

enhanced $I_{Ca,L}$ in a cGMP-independent manner in ferret ventricular myocytes (Campbell *et al.* 1996), but nitrosothiols inhibited the activity of cloned L-type Ca^{2+} channels in HEK293 cells (Hu *et al.* 1997). In rat ventricular myocytes, like in cardiac myocytes from other species and tissues (Méry *et al.* 1993; Levi *et al.* 1994; Kirstein *et al.* 1995; Wahler & Dollinger, 1995; Wang *et al.* 1998; Feron *et al.* 1999), NO-donors (including nitrosothiols and SIN-1) did not elicit a cGMP-independent effect on $I_{Ca,L}$. Interestingly, we found that the basal $I_{Ca,L}$ in rat ventricular myocytes was not modified by reducing agents but was inhibited by LY 83583, a superoxide anion generator. Hu *et al.* (1997) also reported that the basal activity of the L-type Ca^{2+} channel was not changed by reducing agents but was inhibited by oxidant treatments. In contrast, $I_{Ca,L}$ was inhibited by reducing agents (glutathione, DL-DTT) and enhanced by an oxidant (DTNB) in ferret ventricular myocytes (Campbell *et al.* 1996). Altogether, these data suggest that the different effects of NO-donors on the basal $I_{Ca,L}$ might require different redox environments. In addition, it is possible that the “direct” effects of NO-donors do not take place on the L-type Ca^{2+} channel itself, but on other auxiliary proteins and/or on lipids. While the reason for these discrepancies remains to be fully elucidated, our results clearly demonstrate that NO can recruit an endogenous cG-PK in rat ventricular myocytes, leading to a tight control of the β -adrenergic stimulation of L-type Ca^{2+} channel activity.

References

- ABI GERGES, N., ESCHENHAGEN, T., HOVE-MADSEN, L., FISCHMEISTER, R. & MÉRY, P.-F. (1997a). Methylene blue is a muscarinic antagonist in cardiac myocytes. *Molecular Pharmacology* **52**, 482-490.
- ABI GERGES, N., HOVE-MADSEN, L., FISCHMEISTER, R. & MÉRY, P.-F. (1997b). A comparative study of the effects of three guanylyl cyclase inhibitors on the L-type Ca^{2+} and muscarinic K^+ currents in frog cardiac myocytes. *British Journal of Pharmacology* **121**, 1369-1377.
- ABI GERGES, N., MÉRY, P.-F. & FISCHMEISTER, R. (1997c). The NO-sensitive guanylyl cyclase does not participate in the muscarinic regulation of rat cardiac Ca^{2+} current. *Biophysical Journal* **72**, A34.
- ABI-GERGES N., TAVERNIER, B., MEBAZAA, A., FAIVRE, V., PAQUERON, X., PAYEN, D., FISCHMEISTER, R. & MÉRY P.-F. (1999). Sequential changes in autonomic regulation of cardiac myocytes after In vivo endotoxin injection in Rat. *American Journal of Respiratory and Critical Care Medecine* **160**, 1196-1204.
- AKAIKE, T., YOSHIDA, M., MIYAMOTO, Y., SATO, K., KOHNO, M., SASAMOTO, K., MIYAZAKI, K., UEDA, S. & MAEDA, H. (1993). Antagonistic action of imidazolineoxyl N-oxides against endothelium-derived relaxing factor/NO through a radical reaction. *Biochemistry* **32**, 827-832.
- BECKMAN, J. S. & KOPPENOL, W. H. (1996). Nitric oxide, superoxide and peroxynitrite: the good, the bad, and the ugly. *American Journal of Physiology* **271** (Cell Physiol. 40), C1424-C1437.

- BUTT, E., EIGENTHALER, M. & GENEISER H.-G. (1994). (Rp)-8-pCPT-cGMPS, a novel cGMP-dependent protein kinase inhibitor. *European Journal of Pharmacology* **269**, 265-268.
- CAMPBELL, D. L., STAMLER, J. S. & STRAUSS, H. C. (1996). Redox modulation of L-type calcium channels in ferret ventricular myocytes. *Journal of General Physiology* **108**, 277-293.
- FEELISCH, M. (1991). The biochemical pathways of nitric oxide formation from nitrovasodilators: appropriate choice of exogenous NO donors and aspects of preparation and handling of aqueous NO solutions. *Journal of Cardiovascular Pharmacology* **17(suppl.3)**, S25-S33.
- FERON, O., HAN, X. & KELLY, R. A. (1999). Muscarinic cholinergic signaling in cardiac myocytes: dynamic targeting of M2AChR to sarcolemmal caveolae and eNOS activation. *Life Sciences* **64**, 471-477.
- FERRERO, R., RODRIGUEZ-PASCUAL, F., MIRAS-PORTUGAL, M. T. & TORRES, M. (1999). Comparative effects of several nitric oxide donors on intracellular cyclic GMP levels in bovine chromaffin cells: correlation with nitric oxide production. *British Journal of Pharmacology* **127**, 779-787.
- GROSS, W. L., BAK, M. I., INGWALL, J. S., ARSTALL, W. A., SMITH, T. W., BALLIGAND, J.-L. & KELLY, R. A. (1996). Nitric oxide inhibits creatine kinase and regulates heart contractile reserve. *Proceedings of the National Academy of Sciences of the USA* **93**, 5604-5609.
- HILAL-DANDAN, R., URASAWA, K. & BRUNTON, L. L. (1992). Endothelin inhibits adenylate cyclase and stimulates phosphoinositide hydrolysis in adult cardiac myocytes. *Journal of Biological Chemistry* **267**, 10620-10624.
- HU, H., CHIAMVIMONVAT, N., YAMAGISHI, T. & MARBAN, E. (1997). Direct inhibition of expressed cardiac L-type Ca²⁺ channels by S-nitrosothiol nitric oxide donors. *Circulation Research* **81**, 742-752.

ISHIBASHI, T., HAMAGUCHI, M., KATO, K., KAWADA, T., OHTA, H., SASAGE, H. & IMAI, S.

(1993). Relationship between myoglobin contents and increase in cyclic GMP produced by glyceryl trinitrate and nitric oxide in rabbit aorta, right atrium and papillary muscle.

Naunyn-Schmiedeberg's Archiv Pharmacology **347**, 533-561.

KIRSTEIN, M., RIVET-BASTIDE, M., HATEM, S., BÉNARDEAU, A., MERCADIER, J.-J. &

FISCHMEISTER, R. (1995). Nitric oxide regulates the calcium current in isolated human atrial myocytes. *Journal of Clinical Investigation* **95**, 794-802.

KOJDA, G. & KOTTENBERG, K. (1999). Regulation of basal myocardial function by NO.

Cardiovascular Research **41**, 514-523.

KOJDA, G., KOTTENBERG, K., NIX, P., SCHLUTER, K. D., PIPER, H. M. & NOACK, E. (1996).

Low increase in cGMP induced by organic nitrates and nitrovasodilators improves contractile response of rat ventricular myocytes. *Circulation Research* **78**, 91-101.

KOMALAVILA, P. & LINCOLN, T. M. (1996). Phosphorylation of the inositol 1,4,5-triphosphate

receptor. *Journal of Biological Chemistry* **271**, 21933-21938.

LEVI, R. C., ALLOATTI, G., PENNA, C. & GALLO, M. P. (1994). Guanylate-cyclase-mediated

inhibition of cardiac I_{Ca} by carbachol and sodium nitroprusside. *Pflügers Archiv* **426**, 419-426.

MARAGOS, C. M., MORLEY, D., WINK, D. A., DUNAMS, T. M., SAAVEDRA, J. E., HOFFMAN,

A., BOVE, A. A., ISAAC, L., HRABIE, J. A. & KEEFER, L. K. (1991). Complexes of .NO with nucleophiles as agents for the controlled biological release of nitric oxide. Vasorelaxant effects. *Journal of Medical Chemistry* **34**, 3242-3247.

MCDONELL, K. L. & DIAMOND, J. (1997). Cyclic GMP-dependent protein kinase activation in

the absence of negative inotropic effects in the rat ventricle. *British Journal of Pharmacology* **122**, 1425-1435.

- MCDONELL, K. L., TIBBITS, G. F. & DIAMOND, J. (1995). cGMP elevation does not mediate muscarinic agonist-induced negative inotropy in rat ventricular cardiomyocytes. *American Journal of Physiology* **269** (*Heart. Circ. Physiol.* **38**), H1905-H1912.
- MÉRY, P.-F., ABI GERGES, N., VANDECASTEELE, G., JUREVICIUS, J., ESCHENHAGEN, T. & FISCHMEISTER, R. (1997). Muscarinic regulation of the L-type Ca^{2+} current in isolated cardiac myocytes. *Life Sciences* **60**(13/14), 1113-1120.
- MÉRY, P.-F., LOHMANN, S. M., WALTER, U. & FISCHMEISTER, R. (1991). Ca^{2+} current is regulated by cyclic GMP-dependent protein kinase in mammalian cardiac myocytes. *Proceedings of the National Academy of Sciences of the USA* **88**(4), 1197-201.
- MÉRY, P.-F., PAVOINE, C., BELHASSEN, L., PECKER, F. & FISCHMEISTER, R. (1993). Nitric oxide regulates cardiac Ca^{2+} current. Involvement of cGMP-inhibited and cGMP-stimulated phosphodiesterases through guanylyl cyclase activation. *Journal of Biological Chemistry* **268**(35), 26286-95.
- MIYAMOTO, A., LAUFS, U., PARDO, C. & LIAO, J. K. (1997). Modulation of bradykinin receptor ligand binding affinity and its coupled G-proteins by nitric oxide. *Journal of Biological Chemistry* **272**(31), 19601-19608.
- O'BRIEN, P. J., SHEN, H., MCCUTCHEON, L. J., O'GRADY, M., BYRNE, P.J., FERGUSON, H. W., MIRSALEMI, M. S., JULIAN, R. J., SARGEANT, J. M., TREMBLAY, R. R., *et al.* (1992). Rapid, simple and sensitive microassay for skeletal and cardiac muscle myoglobin and hemoglobin: use in various animals indicates functional role of myohemoproteins. *Molecular & Cellular Biochemistry* **112**(1), 45-52.
- PFEIFFER, A., NÜRNBERG, B., KAMM, S., UHDE, M., SCHULTZ, G., RUTH, P. & HOFMANN, F. (1995). Cyclic GMP-dependent protein kinase blocks pertussis toxin-sensitive hormone receptor signaling pathways in chinese hamster ovary cells. *Journal of Biological Chemistry* **270**(16), 9052-9059.

- SANDIRASEGARANE, L. AND DIAMOND, J. (1999). The nitric oxide donors, SNAP and DEA/NO, exert a negative inotropic effect in rat cardiomyocytes which is independent of cyclic GMP elevation. *Journal of Molecular and Cellular Cardiology* **31(4)**, 799-808.
- SCAMPS, F., MAYOUX, E., CHARLEMAGNE, D. & VASSORT, G. (1990). Calcium current in single cells isolated from normal and hypertrophied rat heart. *Circulation Research* **67**, 199-208.
- STEIN, B., DRÖGEMÜLLER, A., MÜLSCH, A., SCHMITZ, W. & SCHOLZ, H. (1993). Ca⁺⁺-dependent constitutive nitric oxide synthase is not involved in the cyclic-GMP-increasing effects of carbachol in ventricular myocytes. *Journal of Pharmacological and Experimental Therapeutics* **266(2)**, 919-925.
- SUMII, K. & SPERELAKIS, N. (1995). cGMP-dependent protein kinase regulation of the L-type Ca²⁺ current in rat ventricular myocytes. *Circulation Research* **77**, 803-812.
- THOMAS, G. P., SIMS, S. M. & KARMAZYN, M. (1997). Differential effects of endothelin-1 on basal and isoprenaline-enhanced Ca²⁺ currents in guinea-pig ventricular myocytes. *Journal of Physiology* **503(1)**, 55-65.
- VALENZUELA, D., HAN, X., MENDE, U., FANKHAUSER, C., MASHIMO, H., HUANG, P., PFEFFER, J., NEER, E. J. & FISHMAN, M. C. (1997). GαO is necessary for muscarinic regulation of Ca²⁺ channels in mouse heart. *Proceedings of the National Academy of Sciences of the USA* **94**, 1727-1732.
- WAHLER, G. M. & DOLLINGER, S. J. (1995). Nitric oxide donor SIN-1 inhibits mammalian cardiac calcium current through cGMP-dependent protein kinase. *American Journal of Physiology* **268** (Cell Physiol. 37), C45-C54.
- WANG, G.-R., ZHU, Y., HALUSHKA, P. V., LINCOLN, T. M. & MENDELSON, M. E. (1998). Mechanism of platelet inhibition by nitric oxide: In vivo phosphorylation of thromboxane

receptor by cyclic GMP-dependent protein kinase. *Proceedings of the National Academy of Sciences of the USA* **95**, 4888-4893.

WANG, Y. G., RECHENMACHER, C. E. & LIPSUS, S. L. (1998). Nitric oxide signaling mediates stimulation of L-type Ca^{2+} current elicited by withdrawal of acetylcholine in cat atrial myocytes. *Journal of General Physiology* **111**, 113-125.

WOLIN, M. S., HINTZE, T. H., SHEN, W., MOHAZZAB-H., K.M. & XIE, Y.-W. (1997). Involvement of reactive oxygen and nitrogen species in signaling mechanisms that control tissue respiration in muscle. *Biochemical Society Transactions* **25**, 934-939.

XU, L., EU, J., MEISSNER, G. & STAMLER, J. (1998). Activation of the cardiac release channel (ryanodine receptor) by poly-S-nitrosylation. *Science* **279**, 234-237.

ZAKHAROV, S. I., PIERAMICI, S., KUMAR, G. K., PRABHAKAR, N. R. & HARVEY, R. D. (1996). Nitric oxide synthase activity in guinea pig ventricular myocytes is not involved in muscarinic inhibition of cAMP-regulated ion channels. *Circulation Research* **78**, 925-935.

Acknowledgements

We thank Patrick Lechêne for skilful technical assistance, Florence Lefebvre for preparation of the cells, Drs. Michel Chesnais and Renzo Levi for helpful discussions. We are grateful to Dr. Xinqiang Han for sharing unpublished results. N. Abi-Gerges was supported by the Fondation pour la Recherche Medicale (F.R.M.) and the Association Française contre les Myopathies (A.F.M.).

Corresponding author

Dr. Najah Abi-Gerges: INSERM U-446, Laboratoire de Cardiologie Cellulaire et Moléculaire, Université Paris-Sud, Faculté de Pharmacie, 5 rue Jean-Baptiste Clément, F-92296 Châtenay-Malabry Cedex, France

Email: Naja.Abi-g@cep.u-psud.fr

Figure Legends

Figure 1. NO-donors do not regulate basal $I_{Ca,L}$ in rat ventricular myocytes

A, a myocyte was first exposed to control intracellular and extracellular control solutions. $I_{Ca,L}$ (\square) was elicited at 0 mV from a holding potential of -50 mV. Superfusion of the myocyte with 100 μ M DEANO is indicated by the horizontal line. Current traces on top were recorded at the times indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. B, Summary of the effects of DEANO (100 μ M), SNP (500 μ M), GSNO (1 mM), SNAP (1 mM), SIN-1 (1 mM) on basal $I_{Ca,L}$ amplitude. The amplitude of $I_{Ca,L}$ in the presence of NO-donors was normalized to the amplitude of basal $I_{Ca,L}$ in control conditions (set to 100%). Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars.

Figure 2. DEANO inhibits the β -adrenergic stimulation of $I_{Ca,L}$ in rat ventricular myocytes

A, a myocyte was first exposed to control intracellular and extracellular solutions. Applications of isoprenaline (Iso 10 nM) and DEANO (100 μ M) are indicated by the horizontal lines. Current traces on top were recorded at the times indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. B (same experiment as in A), Current-voltage relationships of $I_{Ca,L}$ (filled symbols) and of the steady-state current at the end of the pulse (open symbols) obtained under basal conditions, and in the presence of Iso (10 nM) with or without DEANO (100 μ M).

Figure 3. Summary of the effects of DEANO on the β -adrenergic stimulation of $I_{Ca,L}$ in rat ventricular myocytes

The amplitude of $I_{Ca,L}$ in the presence of Iso (1, 3 or 10 nM), without (*empty bars*) or with DEANO (100 μ M, *filled bars*), was normalized to the basal $I_{Ca,L}$ amplitude (set to 100%). Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars. Significant differences from basal (*) or Iso levels (#) are indicated as: * $p < 0.001$; **, ##, $p < 0.0001$.

Figure 4. Other NO-donors do not inhibit the β -adrenergic stimulation of $I_{Ca,L}$ in rat ventricular myocytes

A, a myocyte was first exposed to control intracellular and extracellular solutions. Applications of Iso (10 nM) and SIN-1 (100 μ M) are indicated by the horizontal lines. Current traces on top were recorded at the times indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. B, Summary of the effects of three NO-donors (SNAP, SIN-1, SPNO, at 100 μ M) on $I_{Ca,L}$ in the presence of either 1 nM (*filled bars*) or 10 nM Iso (*empty bars*). The effects of NO-donors on $I_{Ca,L}$ are presented as percent variations from the amplitude of the Iso-stimulated $I_{Ca,L}$. Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars.

Figure 5. The guanylyl cyclase inhibitor ODQ antagonizes the effect of DEANO on the β -adrenergic stimulation of $I_{Ca,L}$ in rat ventricular myocytes

A, a myocyte was first exposed to control intracellular and extracellular solutions. Applications of Iso (3 nM), DEANO (100 μ M) and ODQ (10 μ M) are indicated by the horizontal lines. Current traces on top were recorded at the times indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. B, Summary of the effects of DEANO without (*filled bars*) or with ODQ (10 μ M, *empty bars*), in the presence of 1, 3 or 10 nM Iso. DEANO was used at 10 μ M (in the presence of Iso 10 nM)

or 100 μM (in the presence of 1 and 3 nM Iso). The effects of DEANO on $I_{\text{Ca,L}}$ are presented as percent variations from the amplitude of the Iso-stimulated $I_{\text{Ca,L}}$. Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars. Significant differences from Iso (*) or Iso + DEANO (#) levels are indicated as: *, # $p < 0.05$; ## $p < 0.01$; *** $p < 0.005$.

Figure 6. cG-PK inhibitors antagonize the effect of DEANO on the β -adrenergic stimulation of $I_{\text{Ca,L}}$ in rat ventricular myocytes

A, a myocyte was first exposed to control intracellular and extracellular solutions. Applications of Iso (1 nM), DEANO (100 μM) and Rp8cG (10 μM) are indicated by the horizontal lines. Current traces on top were recorded at the times indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. B, Summary of the effects of DEANO (100 μM) without (*empty bars*) or with cG-PK inhibitors (*filled bars*: Rp8cG 10 μM ; KT5823 0.1 or 0.3 μM), in the presence of 1 or 10 nM Iso. The effects of DEANO on $I_{\text{Ca,L}}$ are presented as percent variations from the amplitude of the Iso-stimulated $I_{\text{Ca,L}}$. Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars. Significant differences from Iso (*) or Iso + DEANO (#) levels are indicated as: *,# $p < 0.05$; ##, $p < 0.01$; ***, $p < 0.001$.

Figure 7. DEANO does not inhibit the stimulation of $I_{\text{Ca,L}}$ induced by exogenous cAMP

A, the pipette solution contained 30 μM cAMP, and cAMP dialysis started when the patch was ruptured (arrow). B, a myocyte was first dialysed with the control solution, and intracellular dialysis with 30 μM cAMP started at the time indicated by the arrow. Applications of DEANO (100 μM) in A or SNP (5, 50 and 500 μM) in B were performed as indicated by the horizontal lines. In A, current traces on top were recorded at the times

indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. *C*, summary of the effects of DEANO, SNP and SNAP on the cAMP (10-100 μM)-stimulated $I_{\text{Ca,L}}$. Variations are given as percent change over the amplitude of the cAMP-stimulated $I_{\text{Ca,L}}$. Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars.

Figure 8. DEANO does not inhibit the β -adrenergic stimulation of $I_{\text{Ca,L}}$ in PTX-treated rat ventricular myocytes

A, a myocyte was incubated with Pertussis toxin (PTX 0.5 $\mu\text{g ml}^{-1}$, 4 hours, 37°C) prior to the experiment. It was first exposed to control extracellular and intracellular solutions, and applications of Iso (1 nM), DEANO (100 μM) and acetylcholine (ACh, 1 μM) were performed as indicated by the horizontal lines. Current traces on top were recorded at the times indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. *B and C*, summary of the effects of DEANO (100 μM) and ACh (1 μM) on the Iso (0.01 and 1 nM)-stimulated $I_{\text{Ca,L}}$ in PTX-treated myocytes. The amplitude of $I_{\text{Ca,L}}$ is presented as percent increase over basal amplitude (in *B*) or as percent variations from the amplitude of the Iso-stimulated $I_{\text{Ca,L}}$ (in *C*). Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars. Significant differences from Iso level are indicated as: *, $p < 0.05$.

Figure 9. Effects of DEANO and ACh on IBMX-stimulated $I_{\text{Ca,L}}$ in rat ventricular myocytes

Untreated (*A*) and PTX-treated (*B*) myocytes were first exposed to control extracellular and intracellular solutions. Applications of IBMX (40 μM in *A*, 20 μM in *B*), DEANO (100 μM , in *A* and *B*) and ACh (1 μM , in *B*) are indicated by the horizontal lines. Current traces on top

were recorded at the times indicated by the corresponding letters on the main graphs. The dotted lines indicate the zero-current level. C, summary of the effects of IBMX (10-80 μM) on $I_{\text{Ca,L}}$, used either alone (*empty bars*) or in the presence of DEANO (100 μM) or ACh (1 μM) (*filled bars*) in untreated (*left part*) or PTX-treated myocytes (*right part*). The effects of IBMX on $I_{\text{Ca,L}}$ are presented as percent variations from the amplitude of basal $I_{\text{Ca,L}}$ (set to 100%). Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars. Significant differences from basal (*) and IBMX (#) levels are indicated as: # $p < 0.05$; **,## $p < 0.01$; ***, $p < 0.005$.

Figure 1. NO-donors do not regulate basal $I_{Ca,L}$ in rat ventricular myocytes

A, a myocyte was first exposed to control intracellular and extracellular control solutions. $I_{Ca,L}$ (□) was elicited at 0 mV from a holding potential of -50 mV. Superfusion of the myocyte with 100 μ M DEANO is indicated by the horizontal line. Current traces on top were recorded at the times indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. B, Summary of the effects of DEANO (100 μ M), SNP (500 μ M), GSNO (1 mM), SNAP (1 mM), SIN-1 (1 mM) on basal $I_{Ca,L}$ amplitude. The amplitude of $I_{Ca,L}$ in the presence of NO-donors was normalized to the amplitude of basal $I_{Ca,L}$ in control conditions (set to 100%). Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars.




Figure 1

Figure 2. DEANO inhibits the β -adrenergic stimulation of $I_{Ca,L}$ in rat ventricular myocytes

A, a myocyte was first exposed to control intracellular and extracellular solutions. Applications of isoprenaline (Iso 10 nM) and DEANO (100 μ M) are indicated by the horizontal lines. Current traces on top were recorded at the times indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. B (*same experiment as in A*), Current-voltage relationships of $I_{Ca,L}$ (*filled symbols*) and of the steady-state current at the end of the pulse (*open symbols*) obtained under basal conditions, and in the presence of Iso (10 nM) with or without DEANO (100 μ M).




Figure 3. Summary of the effects of DEANO on the β -adrenergic stimulation of $I_{Ca,L}$ in rat ventricular myocytes

The amplitude of $I_{Ca,L}$ in the presence of Iso (1, 3 or 10 nM), without (*empty bars*) or with DEANO (100 μ M, *filled bars*), was normalized to the basal $I_{Ca,L}$ amplitude (set to 100%). Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars. Significant differences from basal (*) or Iso levels (#) are indicated as: * $p < 0.001$; **, ##, $p < 0.0001$.




Figure 3

Figure 4. Other NO-donors do not inhibit the β -adrenergic stimulation of $I_{Ca,L}$ in rat ventricular myocytes

A, a myocyte was first exposed to control intracellular and extracellular solutions. Applications of Iso (10 nM) and SIN-1 (100 μ M) are indicated by the horizontal lines. Current traces on top were recorded at the times indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. *B*, Summary of the effects of three NO-donors (SNAP, SIN-1, SPNO, at 100 μ M) on $I_{Ca,L}$ in the presence of either 1 nM (*filled bars*) or 10 nM Iso (*empty bars*). The effects of NO-donors on $I_{Ca,L}$ are presented as percent variations from the amplitude of the Iso-stimulated $I_{Ca,L}$. Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars.




Figure 4

Figure 5. The guanylyl cyclase inhibitor ODQ antagonizes the effect of DEANO on the β -adrenergic stimulation of $I_{Ca,L}$ in rat ventricular myocytes

A, a myocyte was first exposed to control intracellular and extracellular solutions. Applications of Iso (3 nM), DEANO (100 μ M) and ODQ (10 μ M) are indicated by the horizontal lines. Current traces on top were recorded at the times indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. *B*, Summary of the effects of DEANO without (*filled bars*) or with ODQ (10 μ M, *empty bars*), in the presence of 1, 3 or 10 nM Iso. DEANO was used at 10 μ M (in the presence of Iso 10 nM) or 100 μ M (in the presence of 1 and 3 nM Iso). The effects of DEANO on $I_{Ca,L}$ are presented as percent variations from the amplitude of the Iso-stimulated $I_{Ca,L}$. Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars. Significant differences from Iso (*) or Iso + DEANO (#) levels are indicated as: *, # $p < 0.05$; ## $p < 0.01$; *** $p < 0.005$.




Figure 5

Figure 6. cG-PK inhibitors antagonize the effect of DEANO on the β -adrenergic stimulation of $I_{Ca,L}$ in rat ventricular myocytes

A, a myocyte was first exposed to control intracellular and extracellular solutions. Applications of Iso (1 nM), DEANO (100 μ M) and Rp8cG (10 μ M) are indicated by the horizontal lines. Current traces on top were recorded at the times indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. B, Summary of the effects of DEANO (100 μ M) without (*empty bars*) or with cG-PK inhibitors (*filled bars*: Rp8cG 10 μ M; KT5823 0.1 or 0.3 μ M), in the presence of 1 or 10 nM Iso. The effects of DEANO on $I_{Ca,L}$ are presented as percent variations from the amplitude of the Iso-stimulated $I_{Ca,L}$. Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars. Significant differences from Iso (*) or Iso + DEANO (#) levels are indicated as: *,# $p < 0.05$; ##, $p < 0.01$; ***, $p < 0.001$.




Figure 6

Figure 7. DEANO does not inhibit the stimulation of $I_{Ca,L}$ induced by exogenous cAMP

A, the pipette solution contained 30 μ M cAMP, and cAMP dialysis started when the patch was ruptured (arrow). *B*, a myocyte was first dialysed with the control solution, and intracellular dialysis with 30 μ M cAMP started at the time indicated by the arrow. Applications of DEANO (100 μ M) in *A* or SNP (5, 50 and 500 μ M) in *B* were performed as indicated by the horizontal lines. In *A*, current traces on top were recorded at the times indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. *C*, summary of the effects of DEANO, SNP and SNAP on the cAMP (10-100 μ M)-stimulated $I_{Ca,L}$. Variations are given as percent change over the amplitude of the cAMP-stimulated $I_{Ca,L}$. Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars.





Figure 7


Figure 8. DEANO does not inhibit the β -adrenergic stimulation of $I_{Ca,L}$ in PTX-treated rat ventricular myocytes

A, a myocyte was incubated with Pertussis toxin (PTX $0.5 \mu\text{g ml}^{-1}$, 4 hours, 37°C) prior to the experiment. It was first exposed to control extracellular and intracellular solutions, and applications of Iso (1 nM), DEANO (100 μM) and acetylcholine (ACh, 1 μM) were performed as indicated by the horizontal lines. Current traces on top were recorded at the times indicated by the corresponding letters on the main graph. The dotted line indicates the zero-current level. *B and C*, summary of the effects of DEANO (100 μM) and ACh (1 μM) on the Iso (0.01 and 1 nM)-stimulated $I_{Ca,L}$ in PTX-treated myocytes. The amplitude of $I_{Ca,L}$ is presented as percent increase over basal amplitude (in *B*) or as percent variations from the amplitude of the Iso-stimulated $I_{Ca,L}$ (in *C*). Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars. Significant differences from Iso level are indicated as: *, $p < 0.05$.



B PTX-treated myocytes

□ Iso
■ Iso + DEA-NO
□ Iso + ACh



C




Figure 8

Figure 9. Effects of DEANO and ACh on IBMX-stimulated $I_{Ca,L}$ in rat ventricular myocytes

Untreated (A) and PTX-treated (B) myocytes were first exposed to control extracellular and intracellular solutions. Applications of IBMX (40 μ M in A, 20 μ M in B), DEANO (100 μ M, in A and B) and ACh (1 μ M, in B) are indicated by the horizontal lines. Current traces on top were recorded at the times indicated by the corresponding letters on the main graphs. The dotted lines indicate the zero-current level. C, summary of the effects of IBMX (10-80 μ M) on $I_{Ca,L}$, used either alone (*empty bars*) or in the presence of DEANO (100 μ M) or ACh (1 μ M) (*filled bars*) in untreated (*left part*) or PTX-treated myocytes (*right part*). The effects of IBMX on $I_{Ca,L}$ are presented as percent variations from the amplitude of basal $I_{Ca,L}$ (set to 100%). Bars are the means and lines are the S.E.M. of the number of experiments indicated near the bars. Significant differences from basal (*) and IBMX (#) levels are indicated as: # $p < 0.05$; **,## $p < 0.01$; ***, $p < 0.005$.




Figure 9