



**HAL**  
open science

## Secondary B-mode polarization from Faraday rotation in clusters and galaxies

Hiroyuki Tashiro, Nabila Aghanim, Mathieu Langer

► **To cite this version:**

Hiroyuki Tashiro, Nabila Aghanim, Mathieu Langer. Secondary B-mode polarization from Faraday rotation in clusters and galaxies. *Monthly Notices of the Royal Astronomical Society*, 2008, 384 (2), pp.733-746. 10.1111/j.1365-2966.2007.12745.x . hal-03865218

**HAL Id: hal-03865218**

**<https://universite-paris-saclay.hal.science/hal-03865218>**

Submitted on 22 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Secondary $B$ -mode polarization from Faraday rotation in clusters and galaxies

Hiroyuki Tashiro,<sup>★</sup> Nabila Aghanim and Mathieu Langer

*Institut d'Astrophysique Spatiale (IAS), Université Paris-Sud XI and CNRS (UMR 8617), Bâtiment 121, F-91405 Orsay, France*

Accepted 2007 November 20. Received 2007 October 9; in original form 2007 May 20

## ABSTRACT

We revisit the polarization induced by Faraday rotation when cosmic microwave background photons traverse magnetized plasma. We compute the secondary  $B$ -mode angular power spectrum from Faraday rotation due to magnetic fields in galaxies and galaxy clusters with masses ranging from  $10^{11}$  to  $10^{16.5} M_{\odot}$ . We investigate its dependence on the electron and the magnetic field profiles. Namely we consider both the  $\beta$ -profile of electron density as well as an electron density distribution based on the Navarro–Frenk–White dark matter profile. We model the magnetic field structure in galaxies and clusters motivated by recent observations. We further account for its redshift evolution and we examine the importance of its coherence length. We find that the  $B$ -mode polarization from Faraday rotation depends on the normalization parameter  $C_l \propto \sigma_8^5 - \sigma_8^6$ . At 30 GHz for  $\sigma_8 = 0.8$ , the  $B$  modes from Faraday rotation range between 0.01 and  $4 \times 10^{-3} \mu\text{K}^2$  at  $l = 10^4$  in the case of a maximally coherent fields. For smaller coherence lengths, those amplitudes are smaller and they peak at higher multipoles.

**Key words:** magnetic fields – cosmology: theory – large-scale structure of Universe.

## 1 INTRODUCTION

Recent cosmic microwave background (CMB) observations allowed significant progress in cosmology. The CMB temperature anisotropies have been measured in detail and, as a result, we know that the universe is spatially flat and that the power spectrum of the density fluctuations is consistent with scale invariance (Spergel et al. 2006), strongly supporting the inflation scenario.

One of the main targets for the next generation of CMB observations is the detailed measurement of the CMB polarization. The CMB polarization is traditionally split into two components:  $E$  and  $B$  modes (Kamionkowski, Kosowsky & Stebbins 1997b; Zaldarriaga & Seljak 1997). Particularly, the detection of the  $B$ -mode polarization is important for further testing the inflation scenario. Gravitational waves, predicted by inflation scenario but not detected yet, generate the primordial  $B$ -mode polarization, while the density fluctuations do not (Kamionkowski, Kosowsky & Stebbins 1997a; Seljak & Zaldarriaga 1997). Therefore, the measurement of  $B$ -mode polarization is an essential step towards constraining the inflationary scenario.

However, there are other sources of  $B$ -mode polarization. One of them is gravitational lensing (Zaldarriaga & Seljak 1997). Gravitational lensing distorts the CMB polarization fields so that a secondary  $B$ -mode polarization is induced from the primary  $E$ -mode polarization. These secondary  $B$  modes contaminate the primary  $B$ -mode polarization. However, significant effort has been made to find ways how to remove the secondary  $B$ -mode polarization due to gravitational lensing from the polarization maps (see e.g. Hu & Okamoto 2002; Hirata & Seljak 2003), in order to recover the primordial  $B$ -mode angular power spectrum with high accuracy.

The presence of cosmic magnetic fields, observed on various scales, is also an important source for both primary and secondary  $B$ -mode polarization. The main ways of detection and measurement of magnetic fields are Zeeman splitting effect, synchrotron emission and Faraday rotation. Zeeman splitting provides direct observation of magnetic fields. However, the signal is so small with respect to Doppler broadening that it is essentially useless in the cosmological context where magnetic fields are weak and velocities are high. Extragalactic magnetic fields are mainly detected by the other two methods. The amplitude of the fields measured in galaxies and galaxy clusters are typically of the order of 1–10  $\mu\text{G}$ . These methods only provide information on magnetic fields integrated along the line of sight. Nevertheless, in conjunction with complementary measurements (electron number density), they have been successfully applied to show that the coherence length of magnetic fields can be as large as cluster scales, or even bigger (Kim et al. 1989). Similarly, magnetic fields have been measured with the same amplitudes in high-redshift objects at  $z > 2$  (Athreya et al. 1998).

<sup>★</sup>E-mail: hiroyuki.tashiro@ias.u-psud.fr

Although magnetic fields are observed on all scales with increasing accuracy, we still do not know where they originated from. Primordial magnetic fields are one of the candidates for the origin of magnetic fields in galaxies and galaxy clusters. Primordial magnetic fields act as sources of the primary  $B$ -mode polarization by generating vorticity in the cosmic plasma and additional gravitational waves (Seshadri & Subramanian 2001; Mack, Kahniashvili & Kosowsky 2002; Subramanian, Seshadri & Barrow 2003; Lewis 2004; Tashiro, Sugiyama & Banerjee 2006). However, those waves arise on small scales and therefore do not interfere with the detection of the polarization due to inflation-generated gravitational waves.

Magnetic fields also create  $B$ -mode polarization by Faraday rotation, i.e. the rotation of the linear polarization plane through the interaction between CMB photons and magnetized plasma along their path. Faraday rotation distorts the CMB polarization fields like gravitational lensing, so that it induces  $B$ -mode polarization from the  $E$  modes. The effects of Faraday rotation due to primordial magnetic fields were investigated in some papers (Scóccola, Harari & Mollerach 2004; Kosowsky et al. 2005). In addition to primordial magnetic fields, extragalactic magnetic fields of several  $\mu\text{G}$  (Carilli & Taylor 2002) will produce Faraday rotation and therefore add to the  $B$ -mode polarization. This effect was first studied by Takada, Ohno & Sugiyama (2001) who investigated the  $B$ -mode polarization due to Faraday rotation in galaxy clusters. They found that homogeneous magnetic fields of several  $\mu\text{G}$  produce  $1\text{-}\mu\text{K}$   $B$ -mode polarization around  $l = 1000$ . However, it has been shown that magnetic fields in galaxy clusters are not uniform (see Murgia et al. 2004 and references therein). The radial profile of the magnetic fields may affect the  $B$ -mode polarization maps on small scales. Ohno et al. (2003) discuss the possibility of the reconstruction of the magnetic fields in a galaxy cluster from the  $B$ -mode polarization map via the Sunyaev–Zel’dovich (SZ) effect and X-ray emission.

In this paper, we revisit the secondary  $B$ -mode polarization from Faraday rotation in galaxy clusters. We complete the study by taking into account galaxies as well. In spiral galaxies, observations show that the magnetic fields spread widely not only in the galactic plane but also over the halo, with an average amplitude of about  $10\ \mu\text{G}$  (Hummel, Beck & Dahlem 1991). Even in elliptical galaxies, magnetic fields corresponding to many  $\mu\text{G}$  are observed (Greenfield, Roberts & Burke 1985).

The paper is organized as follows. In Section 2, we introduce the formalism of the power spectrum for the Faraday rotation angle and we derive the angular power spectrum of  $B$ -mode polarization due to Faraday rotation. In Section 3, we discuss the two different profiles of gas density and magnetic fields that we use, motivated by observations and numerical simulations. In Section 4, the resulting  $B$ -mode polarization angular power spectrum is shown, taking into account the effects of both galaxies and galaxy clusters. Section 5 is devoted to discussions and summary. Throughout the paper, we use the *Wilkinson Microwave Anisotropy Probe* (WMAP) values of cosmological parameters, i.e.  $h = 0.73$  ( $H_0 = h \times 100\ \text{km s}^{-1}\ \text{Mpc}^{-1}$ ),  $T_0 = 2.725\ \text{K}$ ,  $h^2\ \Omega_b = 0.0223$  and  $h^2\ \Omega_m = 0.128$  (Spergel et al. 2006).

## 2 POLARIZATION ANGULAR POWER SPECTRUM FROM FARADAY ROTATION

### 2.1 Angular power spectrum of the rotation angle

Our aim is to calculate the angular power spectrum of the CMB polarization produced by Faraday rotation on the galaxy and galaxy cluster scales. To do so, we first need to compute the angular power spectrum of the rotation angle.

The rotation angle caused by Faraday rotation is given by

$$\alpha = \frac{2\pi e^3}{m_e^2 \omega^2} \int ds\ n_e B\ \hat{\gamma} \cdot \hat{b}, \quad (1)$$

where  $\omega$  is the angular frequency,  $n_e$  is the free electron number density,  $B$  is the magnetic field strength, and  $\hat{\gamma}$  and  $\hat{b}$  are the directions of the line of sight and the magnetic field, respectively.

We then use the halo formalism commonly used for SZ power spectrum computations (e.g. Cole & Kaiser 1988; Makino & Suto 1993; Komatsu & Kitayama 1999). In this formalism, the angular power spectrum is defined by

$$C_l^\alpha = C_l^{\text{single}} + C_l^{\text{halo-halo}}, \quad (2)$$

where  $C_l^{\text{single}}$  describes the Poisson term and  $C_l^{\text{halo-halo}}$  gives the contribution from correlated haloes. They are expressed as

$$C_l^{\text{single}} = \int_0^{z_{\text{dec}}} dz\ \frac{dV}{dz} \int_{M_{\text{min}}}^{M_{\text{max}}} dM\ \frac{dn(M, z)}{dM} |\alpha_l(M, z)|^2, \quad (3)$$

$$C_l^{\text{halo-halo}} = \int_0^{z_{\text{dec}}} dz\ \frac{dV}{dz} P_m \left( k = \frac{l}{r(z)}, z \right) \left[ \int_{M_{\text{min}}}^{M_{\text{max}}} dM\ \frac{dn(M, z)}{dM} b(M, z) \alpha_l(M, z) \right]^2, \quad (4)$$

where  $r(z)$  and  $V(z)$  are the comoving distance and the comoving volume, respectively. In equations (3) and (4),  $n(M, z)$  is the comoving halo number density of mass  $M$  at redshift  $z$  and  $b(M, z)$  is the linear bias. We adopt the fitting formula given by Sheth & Tormen (1999),

$$n(M, z) dM = \frac{\bar{\rho}}{M} \left[ 1 + 2^{-p} \frac{\Gamma(1/2 - p)}{\sqrt{\pi}} \right] [1 + (qv)^{-p}] \left( \frac{qv}{2\pi} \right)^{1/2} \exp \left( -\frac{qv}{2} \right) \frac{dv}{v}, \quad (5)$$

where  $v = [\delta_c / \sigma(M, z)]^2$ ,  $\delta_c$  is the critical overdensity and  $\sigma$  is the variance smoothed with a top-hat filter of a scale  $R = (3M/4\pi\bar{\rho})^{1/3}$  and we take  $p \approx 0.3$ ,  $q = 0.75$  (Cooray & Sheth 2002). In equation (4), we use the linear bias described as  $b(M, z) = 1 + qv - 1/\delta_c + 2p/\delta_c (1 + (qv)^p)$  (Scoccimarro et al. 2001).

In equations (3) and (4),  $\alpha_l(M, z)$  is the projected Fourier transform of the rotation angle obtained in the small angle approximation,

$$\alpha_l = 2\pi \int d\theta \theta \alpha(\theta, M, z) J_0(l\theta), \quad (6)$$

where  $\alpha(\theta, M, z)$  is the angular profile of the rotation measurement induced by the magnetic field in a galaxy or a galaxy cluster with mass  $M$  at redshift  $z$ . The rotation angle is obtained from equation (1), once a gas distribution and a model for the magnetic field in clusters are given. We set  $\theta = r/D_a$  and  $\theta_c = r_c/D_a$  where  $D_a$  is the angular diameter distance. To compute the power spectrum of the rotation angle, equation (2), we also need to know the angle between the magnetic field and the line of sight. In the following, we assume that the orientation of magnetic fields does not change within a galaxy or a galaxy cluster, and we take  $\langle |\hat{\gamma} \cdot \hat{\mathbf{b}}|^2 \rangle = 1/3$ . This means that the coherence length of magnetic fields is the virial radius and the orientation of the magnetic fields is random from structure to structure. We discuss the case of different coherence lengths in Section 4.2.

## 2.2 B-mode polarization from Faraday rotation

The CMB polarization can be described using the Stokes parameters ( $Q, U$ ). If CMB photons travel in the  $\hat{z}$  direction,  $Q$  is the difference between the intensity in  $\hat{y}$  and  $\hat{x}$  directions, and  $U$  is the difference between intensities in directions obtained by rotating  $\hat{y}$  and  $\hat{x}$  by  $45^\circ$ . If the CMB radiation passes through a magnetized plasma, it undergoes Faraday rotation by an angle  $\alpha$  (equation 1). The Stokes parameters after passing through the plasma can be written as

$$\begin{pmatrix} Q' \\ U' \end{pmatrix} = \begin{pmatrix} \cos 2\alpha & \sin 2\alpha \\ -\sin 2\alpha & \cos 2\alpha \end{pmatrix} \begin{pmatrix} Q \\ U \end{pmatrix}. \quad (7)$$

Under the assumption  $\alpha \ll 1$ , we get

$$\begin{aligned} Q' &\approx Q + 2\alpha U, \\ U' &\approx U - 2\alpha Q. \end{aligned} \quad (8)$$

Accordingly, we can write

$$Q' \pm iU' = (Q + 2\alpha U) \pm i(U - 2\alpha Q) = (1 \mp 2i\alpha)(Q \pm iU). \quad (9)$$

The Stokes parameters depend on the choice of the coordinate system. Therefore it is convenient to introduce the rotation-invariant basis, the  $E$  and  $B$  modes (Kamionkowski et al. 1997b; Zaldarriaga & Seljak 1997). These are obtained by expanding  $Q$  and  $U$  in spin-2 spherical harmonics  ${}_{\pm 2}Y_l^m$ ,

$$Q \pm iU = \sum_{l,m} (E_{lm} \pm iB_{lm}) {}_{\pm 2}Y_l^m. \quad (10)$$

The angle of Faraday rotation can also be decomposed as

$$\alpha = \sum_{l,m} \alpha_{lm} Y_l^m. \quad (11)$$

Applying these decompositions to equation (9), we obtain  $E'_{lm} \pm iB'_{lm}$  after Faraday rotation as

$$\begin{aligned} E'_{lm} \pm iB'_{lm} &= \int d\Omega {}_{\pm 2}Y_l^{m*} \left( 1 \mp 2i \sum_{l_1, m_1} \alpha_{l_1 m_1} Y_{l_1}^{m_1} \right) \sum_{l_2, m_2} (E_{l_2 m_2} \pm iB_{l_2 m_2}) {}_{\pm 2}Y_{l_2}^{m_2} \\ &= E_{lm} \pm iB_{lm} \mp 2i \int d\Omega {}_{\pm 2}Y_l^{m*} \sum_{l_1, m_1} \alpha_{l_1 m_1} Y_{l_1}^{m_1} \sum_{l_2, m_2} (E_{l_2 m_2} \pm iB_{l_2 m_2}) {}_{\pm 2}Y_{l_2}^{m_2}. \end{aligned} \quad (12)$$

From this equation, we can get  $\Delta B = B' - B$ , the contribution of the Faraday rotation to the power spectrum of  $B$  modes. We give the detailed calculation in Appendix A and we only write the result here. The angular power spectrum of the  $B$  mode created by Faraday rotation is written as

$$C_l^{\Delta B} = N_l^2 \sum_{l_1, l_2} N_{l_2}^2 K(l, l_1, l_2)^2 C_{l_2}^E C_{l_1}^\alpha \frac{(2l_1 + 1)(2l_2 + 1)}{4\pi(2l + 1)} (C_{l_1 0 l_2 0}^{l_0})^2, \quad (13)$$

where  $C_{\alpha\beta\gamma}^{c\gamma}$  are the Clebsch–Gordan coefficients,  $N_l = (2(l-2)!/(l+2)!)^{1/2}$  and

$$K(l, l_1, l_2) \equiv -\frac{1}{2} (L^2 + L_1^2 + L_2^2 - 2L_1L_2 - 2L_1L - 2L_2L - 2L), \quad (14)$$

with  $L = l(l+1)$ ,  $L_1 = l_1(l_1+1)$  and  $L_2 = l_2(l_2+1)$ . Pre-existing  $B$ -mode polarization acts like a depolarization source for the  $B$  modes produced by Faraday rotation (see Appendix A, equation A7). In equation (13), we neglect the contributions of such pre-existing  $B$  modes created, for example, by gravitational waves or gravitational lensing. We checked that these contributions are much smaller (two orders magnitude) than primary  $E$ -mode polarization.

### 3 GAS AND MAGNETIC FIELD DISTRIBUTION IN COLLAPSED OBJECTS

In order to evaluate Faraday rotation on the galaxy or galaxy cluster scales, we need the distributions of the electron number density and the magnetic field strength.

#### 3.1 Gas distribution

As for the electron number density, we will consider two cases.

##### 3.1.1 $\beta$ -profile

We first adopt the  $\beta$ -profile (Cavaliere & Fusco-Femiano 1978) for both galaxy cluster and galaxies,

$$n_e(r) = n_c \left( 1 + \frac{r^2}{r_c^2} \right)^{-3\beta/2}, \quad (15)$$

where  $r$ ,  $r_c$  and  $n_c$  are, respectively, the physical distance from the cluster centre, the cluster core radius and the central electron density. We explore a different distribution of the electron number density in Section 3.1.2. Under the assumption of self-similarity  $r_c \propto r_{\text{vir}}$ ,  $r_{\text{vir}}$  being the virial radius. Cluster observations (Mohr, Mathiesen & Evrard 1999) roughly give  $r_{\text{vir}} \approx 10r_c$ . Using the spherical collapse model, the virial radius is related to the mass  $M$  of the dark matter halo and the redshift  $z$  through the following expression:

$$r_{\text{vir}} = \left[ \frac{M}{(4\pi/3)\Delta_c(z)\bar{\rho}(z)} \right]^{1/3}, \quad (16)$$

where  $\Delta_c(z) = 18\pi^2\Omega_m z^{0.427}$  is the spherical overdensity of the virialized halo (Nakamura & Suto 1997).

In the  $\beta$ -profile, the number density of baryons can be written as

$$\rho_b(r) = \mu m_p n_c \left( 1 + \frac{r^2}{r_c^2} \right)^{-3\beta/2}, \quad (17)$$

where  $\mu$  is the mean molecular weight and  $m_p$  is the hydrogen mass. We can safely assume that most of the halo mass is contained within the virial radius. That is,

$$\frac{\Omega_b}{\Omega_m} M = \int_0^{r_{\text{vir}}} dr 4\pi r^2 \rho_b(r). \quad (18)$$

From this equation, we can calculate the central electron density

$$\begin{aligned} n_c &= \frac{\Omega_b}{\Omega_m} \frac{M}{\mu m_p} \left[ \frac{4\pi}{3} r_{\text{vir}}^3 {}_2F_1(3/2, 3\beta/2; 5/2; -(r_{\text{vir}}/r_c)^2) \right]^{-1} \\ &= 9.26 \times 10^{-4} \left( \frac{M}{10^{14} \text{ M}_\odot} \right) \left( \frac{r_{\text{vir}}}{1 \text{ Mpc}} \right)^{-3} \left( \frac{\Omega_b}{\Omega_m} \right) {}_2F_1^{-1}(3/2, 3\beta/2; 5/2; -(r_{\text{vir}}/r_c)^2) \text{ cm}^{-3}, \end{aligned} \quad (19)$$

where  $F_1(\alpha, \beta; \gamma; z)$  is the hypergeometric function. Using  $r_c = 10r_{\text{vir}}$ , we find  ${}_2F_1(3/2, 3\beta/2; 5/2; -(r_{\text{vir}}/r_c)^2)$  is 0.032 for  $\beta = 0.6$  and 0.005 for  $\beta = 1.0$ .

##### 3.1.2 Navarro–Frenk–White profile

We also consider a different electron density profile from the  $\beta$ -profile. Namely we use a density profile motivated by the Navarro–Frenk–White (NFW) dark matter profile (Navarro, Frenk & White 1997).

The NFW dark matter density profile is given by

$$\rho_{\text{dm}}(x) = \frac{\rho_s}{x(1+x)^2}. \quad (20)$$

Here  $x \equiv r/r_s$  where  $r_s$  is a scale radius, and  $\rho_s$  is a scale density at this radius. The scale radius,  $r_s$ , is related to the virial radius

$$r_s(M, z) = \frac{r_{\text{vir}}(M, z)}{c(M, z)}, \quad (21)$$

where  $c$  is the concentration parameter. In the following, we adopt the concentration parameter of Komatsu & Seljak (2002),

$$c \approx \frac{10}{1+z} \left[ \frac{M}{M_*(0)} \right]^{-0.2}, \quad (22)$$

where  $M_*(0)$  is a solution to  $\sigma(M) = \delta_c$  at the redshift  $z = 0$ .

The electron number density profile  $n_e$  can then be obtained from the NFW dark matter profile following Komatsu & Seljak (2002). For this, three assumptions are made: the gas is in hydrostatic equilibrium in the dark matter potential, the gas density follows the dark matter density in the outer parts of the halo and finally the equation of state of the gas is polytropic,  $P_{\text{gas}} \propto \rho_{\text{gas}}^\gamma$  (where  $P_{\text{gas}}$ ,  $\rho_{\text{gas}}$  and  $\gamma$  are the

gas pressure, the gas density and the polytropic index). From these assumptions, with  $n_e \propto \rho_{\text{gas}}/\mu m_p$ , we obtain the electron number density profile

$$n_e(x) = n_c \left\{ 1 - F \left[ 1 - \frac{\ln(1+x)}{x} \right] \right\}^{1/(\gamma-1)}, \quad (23)$$

where the central electron density  $n_c$  and the coefficient  $F$  are

$$n_c = 3.01 \left( \frac{M}{10^{14} M_\odot} \right) \left( \frac{r_{\text{vir}}}{1 \text{ Mpc}} \right)^{-3} \left( \frac{\Omega_b}{\Omega_m} \right) \frac{c}{(1+c)^2} \left[ \ln(1+c) - \frac{c}{1+c} \right]^{-1} \left\{ \frac{c}{c - F[c - \ln(1+c)]} \right\}^{1/(\gamma-1)} \text{ cm}^{-3}, \quad (24)$$

$$F \equiv 3\eta_c^{-1} \frac{\gamma-1}{\gamma} \left[ \frac{\ln(1+c)}{c} - \frac{1}{1+c} \right]^{-1}. \quad (25)$$

Komatsu & Seljak (2002) provide the following useful fitting formulae for  $\gamma$  and  $\eta_c$ :

$$\gamma = 1.137 + 8.94 \times 10^{-2} \ln(c/5) - 3.68 \times 10^{-3}(c-5), \quad (26)$$

$$\eta_c = 2.235 + 0.202(c-5) - 1.16 \times 10^{-3}(c-5)^2. \quad (27)$$

### 3.2 Magnetic field distribution

Magnetic fields in galaxies can reach amplitudes up to 10  $\mu\text{G}$  (Hummel et al. 1991). Observations by Murgia et al. (2004) suggest that the distribution of magnetic fields in galaxies and galaxy clusters is such that their strength decreases outward. Therefore, in the  $\beta$ -profile, we adopt the form proposed by those authors:

$$B(r) = B_c \left( 1 + \frac{r^2}{r_c^2} \right)^{-3\beta\mu/2}, \quad (28)$$

where  $B_c$  is the mean magnetic field strength at the centre,  $\beta$  is the parameter of the  $\beta$ -profile. From equation (15), we get  $B \propto n_e^{\mu\beta}$ . When  $\mu = 1/2$ , the energy density of magnetic fields decreases outward like the electron number density, while in the case where  $\mu = 2/3$ , magnetic fields are frozen into matter.

In the NFW case, the distribution of magnetic fields in galaxies and clusters is given by

$$B(x) = B_c \left\{ 1 - F \left[ 1 - \frac{\ln(1+x)}{x} \right] \right\}^{\mu/(\gamma-1)}. \quad (29)$$

If  $\mu = 2/3$ , magnetic fields are frozen into the matter whereas, if  $\mu = 1/2$ , the energy density of magnetic fields decreases from the centre like the electron number density.

We further model the growth of magnetic fields in galaxies and galaxy clusters, considering the dynamo process which is widely, although not unanimously, accepted (Widrow 2002). In this scenario, magnetic fields are amplified from weak seeds, most likely produced by astrophysical processes, by the dynamo process and the time-scale of the amplification is of the order of the dynamical time-scale,  $t_d = \sqrt{r_{\text{vir}}^3/GM}$ . Assuming typical magnetic field amplitudes of 3 and 10  $\mu\text{G}$  for clusters and galaxies, respectively, we model the magnetic field growth by

$$B_c = \begin{cases} 3 \exp\{-[t_0 - t(z)]/t_d\} \mu\text{G} & 10^{13} \leq M < 10^{16.5} M_\odot \\ 10 \exp\{-[t_0 - t(z)]/t_d\} \mu\text{G} & 10^{11} \leq M < 10^{13} M_\odot \end{cases} \quad (30)$$


where  $t_0$  is the present time.

## 4 RESULTS


In order to evaluate the effect of Faraday rotation due to galaxies and galaxy clusters on CMB polarization, we arbitrarily consider that the contribution from galaxies is associated with haloes of mass lower than  $10^{13} M_\odot$ . Objects with halo masses larger than this limit are considered as galaxy clusters. The particle number density in galaxies is higher than that in galaxy clusters. Moreover, in cold dark matter scenario, the number density of galaxies is higher than that of galaxy clusters. Therefore the Faraday rotation produced in galaxies presumably cannot be neglected against that generated in galaxy clusters.

### 4.1 Rotation angle power spectrum


In Fig. 1, we show the angular power spectrum of the Faraday rotation angle due to galaxy clusters only. We assume that their mass range is between  $10^{13}$  and  $10^{16.5} M_\odot$  and that all clusters follow the  $\beta$ -profile with a magnetic field distributed following equation (28) and  $B_c = 3 \mu\text{G}$ . We checked that the Poisson term in equation (2) dominates in the mass region of galaxy clusters. The contribution from correlated haloes is thus neglected. We note that increasing  $\beta$  increases the amplitude of the power spectrum and shifts its peak to smaller angular scales.



**Figure 1.** The power spectra of Faraday rotation angle caused by 3  $\mu\text{G}$  magnetic fields in galaxy clusters for the  $\beta$ -model case. The solid line is for  $\beta = 0.6$  and  $\mu = 2/3$  and the dashed line is for  $\beta = 0.6$  and  $\mu = 1/2$ . We also plot the case of  $\beta = 1.0$  and  $\mu = 1/2$ , and  $\beta = 1.0$  and  $\mu = 2/3$  as a dotted line and a dash-dotted line, respectively. We use  $\sigma_8 = 0.8$  and set the CMB frequency to 30 GHz.




**Figure 2.** The dependence on the density fluctuation amplitude of the power spectra of Faraday rotation angle. We choose 30 GHz as the CMB frequency and adopt the  $\beta$ -profile with  $\beta = 0.6$ ,  $\mu = 2/3$  and a 3  $\mu\text{G}$  magnetic field. The dashed, the solid, the dotted lines are  $\sigma = 0.9$ , 0.8 and 0.7, respectively.




**Figure 3.** The redshift contribution for various  $l$  modes. The short-dashed, the solid, the long-dashed and the dash-dotted lines represent  $l = 100$ , 500, 1000, 5000 and 10000, respectively. This figure is calculated for the  $\beta$ -profile with  $\beta = 0.6$  and  $\mu = 2/3$  and a 3  $\mu\text{G}$  magnetic field.

This is because large  $\beta$  value gives steeper profiles and smaller core radii. The magnetic field profile also affects the angular power spectrum. However, this dependence is weaker than that of the electron density profile. Fig. 2 shows the dependence of the angular power spectrum on the normalization of the density fluctuation  $\sigma_8$ . We find  $l^2 C_l \propto \sigma_8^5$  somewhat different from the SZ case (Komatsu & Seljak 2002). We compared this result with that of Takada et al. (2001) and found it in general agreement. The location of the peaks in our case and their case is slightly different. This can be explained by the fact that we take into account distribution of the magnetic fields with central peak, while Takada et al. (2001) consider homogeneous magnetic fields over the whole cluster scale.

In Fig. 3, we represent the redshift distribution of  $C_l$ ,  $d \ln C_l / d \ln z$ , for different  $l$  values. The power spectrum of the rotation angle on large scales (small  $l$ ) is mostly due to lower redshift galaxy clusters. The rotation angle strongly depends on the frequency (see equation 1). As a result, low-redshift clusters have a larger contribution to  $C_l$  than in the SZ effect. For example, while most of the contribution at  $l = 10000$  in the SZ power spectrum comes from galaxy clusters at redshift  $z = 2$  (Komatsu & Seljak 2002), the power spectrum of the Faraday rotation angle at the same multipole is associated with galaxy clusters at redshift  $z = 1$ . On scales below  $l = 100$ , the contribution to the rotation angle




**Figure 4.** The mass contribution for various  $l$  modes. The short-dashed, the solid, the long-dashed and the dash-dotted lines represent  $l = 100, 500, 1000, 5000$  and  $10\,000$ , respectively. This figure is calculated for the  $\beta$ -profile with  $\beta = 0.6$  and  $\mu = 2/3$  and a  $3\ \mu\text{G}$  magnetic field.




**Figure 5.** The angular power spectra of Faraday rotation angle caused by galaxy clusters and galaxies. The solid line is the case of the  $\beta$ -profile with  $\beta = 0.6$  and  $\mu = 2/3$  and the dash-dotted line is the case of the  $\beta$ -profile with  $\beta = 0.6$  and  $\mu = 1/2$ . Both are calculated using  $\sigma_8 = 0.8$ . The dashed and the dotted lines are plotted as the case of  $\sigma_8 = 0.9$  and  $0.7$ , respectively. In all plots, the CMB frequency is  $30\ \text{GHz}$ .

$C_l$  from galaxy clusters at redshift lower than  $z = 0.01$  is large. The angular distance of the redshift  $z = 0.01$  is about  $40\ \text{Mpc}$  while the virial radius of the galaxy cluster with mass  $10^{13}\ M_\odot$  is about  $2\ \text{Mpc}$ . This means that the small angle approximation used in equation (6) is not valid at  $l < 100$ . Similarly, we plot the mass distribution of  $C_l$  for different  $l$  values,  $d \ln C_l/d \ln M$ , in Fig. 4. For all  $l$  values, the main contribution is due to galaxy clusters with masses between  $10^{13}$  and  $10^{14}\ M_\odot$ . The amplitude of the rotation angle power spectrum at multipoles larger than  $l = 5000$  is mostly due to clusters with  $10^{13}\ M_\odot$ . Comparing the mass contribution of the SZ effect (Komatsu & Seljak 2002), we find that the Faraday rotation angle is less affected by the massive haloes. This is because the high-redshift clusters do not contribute significantly to the amplitude of the Faraday rotation spectrum (see discussion on the redshift dependence above).


If we take into account the contribution from the whole population of structure including galaxy clusters and galaxies we obtain the results shown in Fig. 5. We calculate both Poisson and halo-halo correlation terms in equation (2) in the mass range  $10^{11} \leq M < 10^{13}\ M_\odot$ . In the mass range of clusters  $10^{13} \leq M < 10^{16.5}\ M_\odot$ , we neglect the halo-halo correlation term as mentioned previously. The peak of the power spectrum is shifted to smaller angular scales ( $l = 20\,000$ ) because of the contribution from galaxies. In Fig. 5, we use  $\beta = 0.6$  and check that varying  $\mu$  in equation (28) does not modify significantly the power spectrum. We also compare the results with different normalization parameters. The dependence of the amplitude of the rotation measurement is approximately  $l^2 C_l \propto \sigma_8^{5.5} - \sigma_8^6$ . The redshift distribution for the power spectrum computed using the  $\beta$ -profile is given in Fig. 6. It also takes into account the evolution of magnetic field equation (30)



**Figure 6.** The redshift contribution to various  $l$  modes. The short-dashed the solid, the long-dashed and the dash-dotted lines represents  $l = 100, 500, 1000, 5000$  and  $10\,000$ , respectively. This figure is calculated in the  $\beta$ -profile with  $\beta = 0.6$  and  $\mu = 2/3$ .



**Figure 7.** The mass contribution to various  $l$  modes. The left-hand panel shows the contribution from the galaxy mass range while the right-hand panel shows that of the galaxy cluster mass range. The short-dashed the solid, the long-dashed and the dash-dotted lines represent  $l = 100, 500, 1000, 5000$  and  $10000$ , respectively. This figure is calculated for the  $\beta$ -profile with  $\beta = 0.6$  and  $\mu = 2/3$ .



**Figure 8.** The angular power spectra of Faraday rotation angle by galaxy clusters and galaxies at 30-GHz CMB frequency. The solid line is the NFW case with magnetic fields frozen into matter,  $\mu = 2/3$ , and the dashed line is the NFW case with magnetic fields which energy density decrease outward like the electron distribution. We also plot the  $\beta$ -profile cases with  $\mu = 2/3$  and  $1/2$  as the dash-dotted and the dotted lines, respectively.

which influences the redshift distribution. The contribution of high-redshift objects (mainly galaxies) is suppressed for  $z > 1$ . Moreover the redshift distribution at high multipoles is shifted to lower  $z$  (see dot-dashed line in Fig. 6). Most of the Faraday rotation-induced signal is thus due to low-redshift objects. If we now explore the mass distribution of  $C_l$  for different  $l$  (Fig. 7), we notice that the contribution from galaxies (left-hand panel) overwhelms that of galaxy clusters (right-hand panel). We also note the difference in amplitudes between the two panels is mainly due to the difference in magnetic field strength between galaxy clusters ( $3 \mu\text{G}$ ) and galaxies ( $10 \mu\text{G}$ ) described in equation (30).

Now, following the same approach as for the  $\beta$ -model, we compute the angular power spectrum of Faraday rotation angle for the NFW profile given in Fig. 8. In this plot, we show the results at 30 GHz, and we assume that the magnetic field distribution within clusters or galaxies writes as equation (29). For comparison, we plot in the same figure the angular power spectrum for the  $\beta$ -profile and magnetic field given by equation (28). The NFW profile is more peaked than the  $\beta$ -profile. As a result, the power spectrum using the NFW profile is shifted to smaller scale (multipoles larger than 30000) as compared to those of  $\beta$ -profile.


#### 4.2 Polarization power spectrum

We show, in Fig. 9, the angular power spectrum of the  $B$ -mode polarization caused by Faraday rotation at 30 GHz using a  $\beta$ -profile for the gas distribution. From equation (13), the  $B$ -mode polarization by Faraday rotation depends on both the primordial  $E$ -mode polarization  $C_l^E$  and the rotation power spectrum  $C_l^r$ . However, since the rotation power spectrum peaks on smaller angular scales than the primordial  $E$ -mode polarization, the produced  $B$ -mode polarization reflects the rotation angle characteristics. They both peak around  $l \sim 8000$  and for a  $\beta$ -profile the peak value as a function of the main parameters of the model is

$$\frac{l(l+1)}{2\pi} C_l^B \approx 3.0 \times 10^{-3} \left( \frac{B_c}{3 \mu\text{G}} \right)^2 \left( \frac{\sigma_8}{0.8} \right)^5 \left( \frac{\nu}{30 \text{ GHz}} \right)^{-4} \mu\text{K}^2. \quad (31)$$

We also give the power spectrum of  $B$  mode from Faraday rotation at 100 GHz (thick lines) which is the optimal band for CMB observation.

For comparison, we plot in the same figure the power spectrum of the primary  $E$ -mode polarization (thin dotted line), the  $B$ -mode polarization due to the gravitational waves (thin dot-dashed line) and the lensing  $B$ -mode polarization (thin dot-dotted line). The polarization from gravitational waves is computed with a tensor to scalar ratio  $r = 0.6$  which is the upper limit from the *WMAP* three years data (Spergel et al. 2006). This signal dominates on large angular scales. On small scales, the power spectrum of the polarization caused by Faraday rotation



**Figure 9.** The angular power spectra of the  $B$ -mode CMB polarization caused by Faraday rotation for the  $\beta$ -model. The solid line is for  $\beta = 0.6$  and  $\mu = 2/3$  the dashed line is for  $\beta = 0.6$  and  $\mu = 1/2$ , the dash-dotted line  $\beta = 0.6$  and  $\mu = 2/3$  and the dotted line is for  $\beta = 1.0$  and  $\mu = 1/2$ . We use  $\sigma_8 = 0.8$  and set the CMB frequency to 30 GHz. For comparison, we plot, for 100 GHz, the spectrum for  $\beta = 0.6$  and  $\mu = 2/3$  (thick solid line), and for  $\beta = 0.6$  and  $\mu = 2/3$  (thick dashed line). We also give the power spectra of the primordial  $E$ -mode polarization, the  $B$ -mode polarization caused by gravitational waves with a tensor to scalar ratio  $r = 0.6$ , and the gravitational lensing  $B$ -mode polarization.

dominates the  $B$ -mode polarization due to gravitational lensing at frequencies lower than 30 GHz, for reasonable values of the cluster magnetic field. When the frequency increases, the gravitational lensing  $B$ -mode polarization dominates. We plot the power spectrum of the Faraday induced polarization for different parameters  $\beta$  and  $\mu$  describing the electron density and the magnetic field profiles (see line styles in caption). Similarly to the rotation angle power spectrum Fig. 1, we note that varying  $\mu$  has very little effect whereas varying  $\beta$  changes the amplitude of the power spectrum by two orders of magnitude.

In the above calculation, we assumed that the coherence length of the magnetic field in a galaxy cluster is the virial radius. This means that the inner product of the directions of the line of sight and of the magnetic field in equation (1) is constant over a galaxy cluster. However, many observations suggest that the coherence length is shorter than the virial radius (e.g. Brandenburg & Subramanian 2005). In order to evaluate the effect of different coherence lengths, we assume that the direction of the magnetic field in a galaxy cluster changes smoothly and the inner product of directions in equation (1) has the following form:


$$\hat{\gamma} \cdot \hat{\mathbf{b}} = \cos\left(\frac{s}{l_c} \pi\right), \quad (32)$$

where  $l_c$  is the coherence length. Taking  $l_c = r_{\text{vir}}/3$ ,  $r_{\text{vir}}/4$  and  $r_{\text{vir}}/10$ , we calculate the angular power spectrum of  $B$ -mode polarization and plot the results in Fig. 10. The small coherence length implies many changes of the direction of magnetic fields so that depolarization occurs along the line of sight. When the coherence length is one-tenth of the virial radius, the amplitude of the angular spectrum is decreased and the peak position shifts to small scales.


Next, the angular power spectrum of  $B$ -polarization caused by Faraday rotation in galaxy clusters and galaxies is plotted in Fig. 11. We plot this figure for a  $\beta$ -profile with  $\beta = 0.6$  and  $\mu = 2/3$ , and at a frequency of 30 GHz. The power spectrum peaks around  $l = 10\,000$  with a peak amplitude, which depends on  $\sigma_8$ ,  $\nu$  and  $B_c$ ,

$$\frac{l(l+1)}{2\pi} C_l^B \approx 1.1 \times 10^{-2} \left(\frac{B_c}{10 \mu\text{G}}\right)^2 \left(\frac{\sigma_8}{0.8}\right)^6 \left(\frac{\nu}{30 \text{GHz}}\right)^{-4} \mu\text{K}^2. \quad (33)$$

The  $B$ -mode polarization induced by galaxies can be the major component of  $B$ -mode polarization on small scales. If the average magnetic field of galaxies is 10  $\mu\text{G}$ , the Faraday rotation produces the dominant  $B$ -mode polarization at 30 GHz for  $l > 4000$ . Alternatively, on scales



**Figure 10.** The angular power spectra of the  $B$ -mode CMB polarization by Faraday rotation for the different coherence lengths. The solid line corresponds to the model with  $l_c = r_{\text{vir}}$ . The dashed, the dash-dotted and the dotted lines are for  $l_c = r_{\text{vir}}/3$ ,  $l_c = r_{\text{vir}}/4$  and  $l_c = r_{\text{vir}}/10$ . For comparison, we plot the  $B$ -mode polarization caused by gravitational waves.



**Figure 11.** The angular power spectra of the  $B$ -mode polarization caused by Faraday rotation galaxy clusters and galaxies for the  $\beta$ -profile. The solid line is for  $\beta = 0.6$  and  $\mu = 2/3$ , and the dash-dotted line is for  $\beta = 0.6$  and  $\mu = 1/2$ . We also plot the case of  $\sigma_8 = 0.9$  and  $0.7$  for  $\beta = 0.6$  and  $\mu = 2/3$  (dashed and dotted lines, respectively). For comparison, we give the  $B$ -mode polarization induced by gravitational waves with  $r = 0.6$ .

smaller than  $l = 10000$ , the  $B$ -mode polarization generated by galaxies for the same magnetic field is expected to dominate at frequencies up to 60 GHz.

As a galaxy evolves, gas is falling from the halo on to the disc and the halo gas density decreases. The remaining fraction of diffuse baryons in the galaxy halo at low redshift can be as low as 10 per cent, whereas in a galaxy cluster it roughly stays equal to 1 (Binney & Merrifield 1998). The evolution of the gas in a galaxy is expected to affect the power spectrum. We estimate the effect of the gas depletion by assuming that only one-tenth of the total baryon density is left in the halo and the rest is condensated on the galactic disc, so that

$$\alpha = \frac{2\pi e^3}{m_e^2 \omega^2} \int ds f_g n_e B \hat{\gamma} \cdot \hat{b}, \quad (34)$$


where

$$f_g = \begin{cases} 0.1, & M < 10^{14} M_\odot, \\ 1, & M > 10^{14} M_\odot. \end{cases} \quad (35)$$


90 per cent depletion is the maximum we can expect. Therefore, results in the left-hand panel of Fig. 12 corresponding to equation (35) show the maximum impact of gas depletion on the power spectrum of  $B$ -mode polarization.

The contribution from galaxies appears only at small scales and is totally overwhelmed by that from galaxy clusters. This result tells us that we may neglect Faraday rotation from galaxies, if most of gas is condensed on to the disc.

In Fig. 11, we showed the  $B$ -mode power spectrum taking into account magnetic field evolution according to equation (30). However, the magnetic field evolution, especially in galaxy clusters, is not fully understood yet. In some scenarios, the time-scale of the evolution is shorter than the dynamical time-scale (e.g. Brandenburg & Subramanian 2005). To highlight the importance of the magnetic field evolution, we calculate the angular spectrum in the case of constant magnetic fields and we plot the results in the right-hand panel in Fig. 12. For higher  $l$  modes, the main contribution comes from galaxies at redshifts higher than  $z = 1$ . Due to this, in the case of constant magnetic fields, the amplitude of the power spectrum is higher and the tail at high  $l$  is also amplified.



**Figure 12.** Left-hand panel: The angular power spectra of the  $B$ -mode polarization caused by Faraday rotation in galaxy clusters and galaxies. For the solid line, we assume that 90 per cent of gas in galaxies is condensed in the disc. The dotted line shows the contribution from galaxy clusters only. For comparison, we plot the power spectrum for the case without gas depletion and that due to gravitational waves with  $r = 0.6$  as the dashed and the dash-dotted lines, respectively. Right-hand panel: The angular power spectra of the  $B$ -mode polarization for different evolutions of magnetic fields. The solid line is for the model where magnetic fields are constant. The dotted line corresponds to the case where magnetic fields are constant and gas is depleted in galactic haloes. We also plot the case where the time-scale of the magnetic field evolution is the dynamical time-scale as the dashed line. The dash-dotted line is taken into account both magnetic field evolution and gas depletion.



**Figure 13.** The angular power spectra of the  $B$ -mode polarization caused by Faraday rotation from galaxy clusters and galaxies at 30 GHz. The solid line is the case of the NFW profile with  $\mu = 2/3$ , and the dashed line is for  $\mu = 1/2$ . For comparison, we plot the power spectra for  $\beta$ -profiles with  $\beta = 0.6$  and  $\mu = 2/3$  and  $\beta = 0.6$  and  $\mu = 1/2$  (dash-dotted and dotted lines, respectively). We also show the  $B$ -mode polarization induced by gravitational waves with  $r = 0.6$ .

Finally, we plot the angular power spectrum of the  $B$ -mode polarization induced by Faraday rotation from galaxy clusters and galaxies with the NFW profile in Fig. 13. Here we neglect the gas depletion in galaxies and we consider the magnetic field evolution, equation (30). Using the NFW profile shifts the polarization spectrum to smaller scales. For  $\mu = 2/3$  the spectrum peaks at  $l > 20\,000$ , and the value at  $l = 10\,000$  is about

$$\frac{l(l+1)}{2\pi} C_l^B \approx 4.0 \times 10^{-3} \left( \frac{B_c}{10 \mu\text{G}} \right)^2 \left( \frac{\nu}{30 \text{ GHz}} \right)^{-4} \mu\text{K}^2. \quad (36)$$

## 5 DISCUSSION AND SUMMARY

In this study, we calculated the secondary  $B$ -mode polarization caused by Faraday rotation. Particularly, we investigated the dependence of the  $B$ -mode angular power spectrum from Faraday rotation on the electron and the magnetic field profiles in galaxies and galaxy clusters. We considered both the  $\beta$ -profile of electron density as well as an electron density distribution based on the NFW dark matter profile. We modelled the magnetic field structure in galaxies or galaxy clusters motivated by observations and further accounted for the redshift evolution of its amplitude.

We showed that the electron and magnetic field profiles in galaxies and galaxy clusters modify the position of the spectrum peak. The amplitude of magnetic fields affects the amplitude of the spectrum, mainly. The  $B$ -mode polarization from Faraday rotation at 30 GHz is  $0.01 \mu\text{K}^2$  at  $l = 10^4$  for the  $\beta$ -profile, whereas it is  $4 \times 10^{-3} \mu\text{K}^2$  at the same  $l$  for the NFW profile. We also investigated the impact of different coherence lengths on the angular power spectrum. Small coherence length induces the depolarization and the peak of the power spectrum shifts to smaller scales and its amplitude is suppressed.

The Faraday rotation angle power spectrum as well as the  $B$ -mode angular spectrum are dominated by the contribution from galaxies at redshifts  $z < 1$  and with masses from  $10^{11}$  to  $10^{12} M_\odot$ . In other words, the detection of a secondary polarization from Faraday rotation would give the average strength of magnetic fields in low-redshift galaxies. However, the  $B$ -mode angular power spectrum is strongly dependent on the evolution of magnetic fields in galaxies and galaxy clusters, as well as on the diffuse gas fraction in galactic haloes. If the time-scale of the magnetic field evolution is shorter than the dynamical time-scale, the contribution from galaxies at redshifts higher than  $z = 1$  is dominant. The power spectrum has higher amplitude and the peak is on smaller scales. Conversely, if the gas evolution is rapid and most of gas in galaxies condensed on to their discs, the contribution from galaxies becomes much smaller and can be neglected.

The contribution from Faraday rotation from clusters and galaxies appears always small enough, on large scales, to be neglected as compared with the primary  $B$  modes from gravitational waves. It is also important to compare the secondary  $B$  modes from Faraday rotation with the secondary  $B$  modes caused by gravitational lensing. The  $B$ -mode polarization by Faraday rotation dominates that due to gravitational lensing at  $l > 8000$  at 30 GHz. It has been shown that gravitational lensing polarization can be removed from the CMB polarization map leaving a residual contribution of about only 10 per cent (Hu & Okamoto 2002; Hirata & Seljak 2003). Considering this cleaning, the  $B$ -mode polarization from Faraday rotation will exceed the remaining signal from lensing at  $l = 2000$  (30 GHz), or at  $l = 7000$  (100 GHz). These values are obtained in the  $\beta$ -profile case with  $\sigma_8 = 0.8$ .

There are other contributions from secondary polarization sources on small scales that we compare to our results. One of them is polarization from inhomogeneous reionization. The coupling between the primordial CMB temperature quadrupole anisotropy and the density fluctuations of free electrons due to inhomogeneous reionization produces a second-order polarization. Some authors estimated this polarization by using a semi-analytic approach (Liu et al. 2001; Mortonson & Hu 2006). For example, Liu et al. (2001) gave  $l(l+1)C_l^B/2\pi \approx 10^{-5} \mu\text{K}^2$  at  $l \sim 10^4$ . Recently, Doré et al. (2007) have revisited this topic using up-to-date numerical simulation. Their results are consistent with those of Liu et al. (2001). The polarization from Faraday rotation due to the realistic magnetic fields in galaxies and galaxy clusters thus dominates the secondary polarization from inhomogeneous reionization, at 30 GHz. However, Faraday rotation has a strong dependence on frequency,

$C_l \propto \nu^4$ . As a result, the polarization by Faraday rotation becomes subdominant with respect to the polarization due to inhomogeneous reionization when the frequency exceeds 60 GHz. Next, we considered the polarization from ionized gas in filaments and large-scale structure (LSS). The primary quadrupole produces the polarization by interaction with the free electron gas captured in potential wells of LSS. Liu, da Silva & Aghanim (2005) have evaluated the resulting polarization and obtained  $l(l+1)C_l^B/2\pi \approx 10^{-3} \mu\text{K}^2$  at  $l \sim 10^4$ . This value is somewhat lower than the polarization due to Faraday rotation at 30 GHz. While the Faraday rotation depends on frequency, the polarization from LSS does not. Therefore, although the polarization of the Faraday rotation dominates the polarization of LSS at frequencies lower than 30 GHz, the former is subdominant for  $\nu < 30$  GHz.

Another source of polarization that needs to be considered is galactic foregrounds. On small angular scales, the main contributions come from synchrotron emission and dust in our Galaxy. Tucci et al. (2005) estimated these contributions, focusing on the polarization at multipoles larger than  $l = 1000$ . We extrapolated their results to smaller scales and obtained, straightforwardly,  $l(l+1)C_l^B/2\pi \approx 1 \mu\text{K}^2$  at  $l \sim 10^4$  at 30 GHz. In order to detect the polarization caused by Faraday rotation, we need to remove the galactic foreground polarization carefully as it will be a major contamination. However, we have to bear in mind that such an extrapolation relies on the assumption that there is significant dust contribution on scales,  $l \approx 10^4$ . On one hand, present observations indicate that the power spectrum of the dust fluctuations decrease as fast as  $k^{-3}$  (Miville-Deschênes, Lagache & Puget 2002) suggesting a fast decrease of the dust contribution on small scales. On the other hand, the contribution from synchrotron emission on those small scales depends on the structure and amplitude of the galactic magnetic field on the same scales. The distribution of the magnetic fields in the Galaxy has been widely studied. Magnetic fields in the Galaxy will, therefore, also act as a Faraday rotation source (Dineen & Coles 2005), but their contributions need to be studied in more detail.

## ACKNOWLEDGMENT

We thank an anonymous referee for useful comments to improve our paper.

## REFERENCES

- Athreya R. M., Kapahi B. N., McCarthy P. J., van Breugel W., 1998, *A&A*, 329, 809  
 Binney J. J., Merrifield M., 1998, *Galactic Astronomy*, 1st edn. Princeton Univ. Press, Princeton, NJ  
 Brandenburg A., Subramanian K., 2005, *Phys. Rep.*, 417, 1  
 Carilli C. L., Taylor G. B., 2002, *ARA&A*, 40, 319  
 Cavaliere A., Fusco-Femiano R., 1978, *A&A*, 70, 677  
 Cole S., Kaiser N., 1988, *MNRAS*, 233, 637  
 Cooray A., Sheth R., 2002, *Phys. Rep.*, 372, 1  
 Dineen P., Coles P., 2005, *MNRAS*, 362, 403  
 Doré O., Holder G., Alvarez M., Iliev I. T., Mellema G., Pen U.-L., Shapiro P. R., 2007, *Phys. Rev. D*, 76, 043002  
 Greenfield P. D., Roberts D. H., Burke B. F., 1985, *ApJ*, 293, 370  
 Hirata C. M., Seljak U., 2003, *Phys. Rev. D*, 67, 043001  
 Hu W., Okamoto T., 2002, *ApJ*, 574, 566  
 Hummel E., Beck R., Dahlem M., 1991, *A&A*, 248, 23  
 Kamionkowski M., Kosowsky A., Stebbins A., 1997a, *Phys. Rev. Lett.*, 78, 2058  
 Kamionkowski M., Kosowsky A., Stebbins A., 1997b, *Phys. Rev. D*, 55, 7368  
 Kim K.-T., Kromberg P. P., Giovannini G., Venturi T., 1989, *Nat*, 341, 720  
 Komatsu E., Kitayama T., 1999, *ApJ*, 526, L1  
 Komatsu E., Seljak U., 2002, *MNRAS*, 336, 1256  
 Kosowsky A., Kahnishvili T., Lavrelashvili G., Ratra B., 2005, *Phys. Rev. D*, 71, 043006  
 Lewis A., 2004, *Phys. Rev. D*, 70, 043011  
 Liu G.-C., Sugiyama N., Benson A. J., Lacey C. G., Nusser A., 2001, *ApJ*, 561, 504  
 Liu G.-C., da Silva A., Aghanim N., 2005, *ApJ*, 621, 15  
 Mack A., Kahnishvili T., Kosowsky A., 2002, *Phys. Rev. D*, 65, 123004  
 Makino N., Suto Y., 1993, *ApJ*, 405, 1  
 Miville-Deschênes M.-A., Lagache G., Puget J.-L., 2002, *A&A*, 393, 749  
 Mohr J. J., Mathiesen B., Evrard A. E., 1999, *ApJ*, 517, 627  
 Mortonson M. J., Hu W., 2007, *ApJ*, 657, 1  
 Murgia M., Govoni F., Feretti L., Giovannini G., Dallacasa D., Fanti R., Taylor G. B., Dolag K., 2004, *A&A*, 424, 429  
 Nakamura T. T., Suto Y., 1997, *Prog. Theor. Phys.*, 97, 49  
 Navarro J. F., Frenk C. S., White S. D. M., 1997, *ApJ*, 490, 493  
 Ohno H., Takada M., Dolag K., Bartelmann M., Sugiyama N., 2003, *ApJ*, 584, 599  
 Page L. et al., 2007, *ApJS*, 170, 335  
 Scocimarro R., Sheth R. K., Hui L., Jain B., 2001, *ApJ*, 546, 20  
 Scóccola C., Harari D., Mollerach S., 2004, *Phys. Rev. D*, 70, 063003  
 Seljak U., Zaldarriaga M., 1997, *Phys. Rev. Lett.*, 78, 2054  
 Seshadri T. R., Subramanian K., 2001, *Phys. Rev. Lett.*, 87, 101301  
 Sheth R. K., Tormen G., 1999, *MNRAS*, 308, 119  
 Spergel D. N. et al., 2007, *ApJS*, 170, 377  
 Subramanian K., Seshadri T. R., Barrow J. D., 2003, *MNRAS*, 344, L31  
 Takada M., Ohno H., Sugiyama N., 2001, preprint (astro-ph/0112412)

Tashiro H., Sugiyama N., Banerjee R., 2006, Phys. Rev. D, 73, 023002  
 Tucci M., Martínez-González E., Vielva P., Delabrouille J., 2005, MNRAS, 360, 935  
 Varshalovich D. A., Moskalev A. N., Khersonskii V. K., 1988, Quantum Theory of Angular Momentum. World Scientific Press, Singapore  
 Widrow L. M., 2002, Rev. Mod. Phys., 74, 775  
 Zaldarriaga M., Seljak U., 1997, Phys. Rev. D, 55, 1830

## APPENDIX A: B-MODE POWER SPECTRUM CALCULATIONS

In this appendix, we deal with the derivation of equation (13) from equation (12). Equation (12) involves the integrations of the spin-weighted spherical harmonics in the term introduced by Faraday rotation. The integrations of the spin-weighted spherical harmonics are calculated through the Clebsch–Gordan coefficients,

$$\begin{aligned} \int d\Omega {}_{\pm 2}Y_l^{m*} Y_{l_1}^{m_1} {}_{\pm 2}Y_{l_2}^{m_2} &= \int d\Omega {}_{\pm 2}Y_l^{m*} \sqrt{\frac{2l_1+1}{4\pi}} \sqrt{\frac{2l_2+1}{4\pi}} \sum_{l_3 m_3} C_{l_1 m_1 l_2 m_2}^{l_3 m_3} C_{l_1 0 l_2 \mp 2}^{l_3 \mp 2} \sqrt{\frac{4\pi}{2l_3+1}} {}_{\pm 2}Y_{l_3}^{m_3} \\ &= \sqrt{\frac{2l_1+1}{4\pi}} \sqrt{\frac{2l_2+1}{2l+1}} C_{l_1 m_1 l_2 m_2}^{lm} C_{l_1 0 l_2 \mp 2}^{l \mp 2}, \end{aligned} \quad (\text{A1})$$

where we use the following property of the spin weighted harmonics:

$${}_s Y_{l_1}^{m_1} {}_{s_2} Y_{l_2}^{m_2} = \sqrt{\frac{2l_1+1}{4\pi}} \sqrt{\frac{2l_2+1}{4\pi}} \sum_{l_3 m_3 s_3} C_{l_1 m_1 l_2 m_2}^{l_3 m_3} C_{l_1 -s_1 l_2 -s_2}^{l_3 -s_3} \sqrt{\frac{4\pi}{2l_3+1}} {}_{s_3} Y_{l_3}^{m_3} \quad (\text{A2})$$

and the orthogonality condition

$$\int d\Omega {}_s Y_{l_1}^{m_1*} {}_{s'} Y_{l_2}^{m_2} = \delta_{m_1 m_2} \delta_{l_1 l_2}. \quad (\text{A3})$$

Substituting equation (A1) to equation (12), we obtain the representation of  $B$ -mode polarization from Faraday rotation

$$B'_{lm} = B_{lm} - \sum_{l_1 m_1} \sum_{l_2 m_2} [1 + (-1)^{l_1+l_2-l}] \sqrt{\frac{2l_1+1}{4\pi}} \sqrt{\frac{2l_2+1}{2l+1}} C_{l_1 m_1 l_2 m_2}^{lm} C_{l_1 0 l_2 2}^{l 2} \alpha_{l_1 m_1} (E_{l_2 m_2} - i B_{l_2 m_2}), \quad (\text{A4})$$

where we use the following relation to obtain the last equation:

$$C_{a\alpha b\beta}^{c\gamma} = (-1)^{a+b-c} C_{a-\alpha b-\beta}^{c-\gamma}. \quad (\text{A5})$$

Next, we derive the  $B$ -mode polarization produced by Faraday rotation  $\Delta B_{lm}$ . We assume that the rotation fields and the primordial polarization are statistically isotropic and uncorrelated,

$$\langle \alpha_{l_1 m_1}^* E_{l_2 m_2}^* \alpha_{l_3 m_3} E_{l_4 m_4} \rangle = \delta_{l_1 l_3} \delta_{m_1 m_3} \delta_{l_2 l_4} \delta_{m_2 m_4} C^\alpha(l_1) C^E(l_2). \quad (\text{A6})$$

The angular power spectrum of the  $B$ -mode polarization from Faraday rotation  $C^{\Delta B}(l)$  is given by

$$\begin{aligned} C_l^{\Delta B} &= \langle |\Delta B_{lm}|^2 \rangle \\ &= \sum_{l_1} \sum_{l_2} [(1 + (-1)^{l_1+l_2-l}) C_{l_1 0 l_2 2}^{l 2}]^2 \frac{2l_1+1}{4\pi} \frac{2l_2+1}{2l+1} C_{l_1}^\alpha (C_{l_2}^E - C_{l_2}^B), \end{aligned} \quad (\text{A7})$$

where we use the orthogonality of the Clebsch–Gordan coefficients,  $\sum_{\alpha\beta} C_{a\alpha b\beta}^{c\gamma} C_{a\alpha b\beta}^{c'\gamma'} = \delta_{cc'} \delta_{\gamma\gamma'}$ . Equation (A7) shows that the Faraday rotation produces  $B$ -mode polarization from  $E$ -mode polarization and transforms a part of the pre-existing  $B$ -mode polarization into  $E$  modes.

It is hard to calculate equation (A7) directly. To reduce the amount of the calculations, we rewrite equation (A7) using the following equations about the Clebsch–Gordan coefficients (Varshalovich, Moskalev & Khersonskii 2005),

$$C_{a\alpha b\beta}^{c\gamma} = (-1)^{a-\alpha} \sqrt{\frac{2c+1}{2b+1}} C_{c\beta a-\alpha}^{b\gamma} = (-1)^{b+\beta} \sqrt{\frac{2c+1}{2a+1}} C_{c-\gamma b\beta}^{a-\alpha} = (-1)^{b+\beta} \sqrt{\frac{2c+1}{2a+1}} C_{b-\beta c\gamma}^{a\alpha}, \quad (\text{A8})$$

$$C_{a\alpha b\beta}^{c\gamma \mp 1} = \sqrt{\frac{(a \mp \alpha)(a \pm \alpha + 1)}{(c \pm \gamma)(c \mp \gamma + 1)}} C_{a\alpha \pm 1 b\beta}^{c\gamma \mp 1} + \sqrt{\frac{(b \mp \beta)(b \pm \beta + 1)}{(c \pm \gamma)(c \mp \gamma + 1)}} C_{a\alpha b\beta \pm 1}^{c\gamma \mp 1}, \quad (\text{A9})$$

$$C_{a1 b-1}^{c0} = \sqrt{\frac{(c+1) - a(a+1) - b(b+1)}{2\sqrt{a(a+1)b(b+1)}}} C_{a0 b0}^{c0}, \quad (\text{A10})$$

$$C_{a1 b1}^{c2} = \frac{a(a+1)[c(c+1) - a(a+1) + b(b+1)] + b(b+1)[c(c+1) + a(a+1) - b(b+1)]}{2\sqrt{a(a+1)b(b+1)c(c-1)(c+1)(c+2)}} C_{a0 b0}^{c0}. \quad (\text{A11})$$

After a lengthy computation, we obtain

$$C_l^{\Delta B} = N_l^2 \sum_{l_1 l_2} N_{l_2}^2 K(l, l_1, l_2)^2 C_{l_1}^\alpha \frac{(2l_1+1)(2l_2+1)}{4\pi(2l+1)} (C_{l_1 0 l_2 0}^{l 0})^2 (C_{l_2}^E - C_{l_2}^B), \quad (\text{A12})$$

where

$$N_l = [2(l-2)!/(l+2)!]^{1/2} \quad (\text{A13})$$

and

$$K(l, l_1, l_2) \equiv -\frac{1}{2} (L^2 + L_1^2 + L_2^2 - 2L_1L_2 - 2L_1L + 2L_1 - 2L_2 - 2L), \quad (\text{A14})$$

with  $L = l(l+1)$ ,  $L_1 = l_1(l_1+1)$  and  $L_2 = l_2(l_2+1)$ .

To compute the Clebsch–Gordan coefficients in equation (13), we utilize the approximation of Kosowsky et al. (2005):

$$\frac{(C_{a0b0}^c)^2}{2c+1} \approx \frac{e}{2\pi} \left(1 + \frac{1}{2g}\right)^{-2g-3/2} \exp \left[ \frac{1}{8g} - \frac{1}{8(g-a)} - \frac{1}{8(g-b)} - \frac{1}{8(g-c)} \right] [g(g-a)(g-b)(g-c)]^{-1/2}. \quad (\text{A15})$$

This approximation is accurate to only 1 per cent for the worst case  $a = b = c = 2$ .

This paper has been typeset from a  $\text{\TeX/L\AA\TeX}$  file prepared by the author.